arXiv:0706.2229v2 [math.OA] 17 Jun 2007

# AF-embedding of the crossed products of AH-algebras by finitely generated abelian groups

Huaxin Lin

**Abstract**

Let $X$ be a compact metric space and let $\Lambda$ be a $\mathbb{Z}^k$ ($k \geq 1$) action on $X$. We give a solution to a version of Voiculescu's problem of AF-embedding: The crossed product $C(X) \rtimes_\Lambda \mathbb{Z}^k$ can be embedded into a unital simple AF-algebra if and only if $X$ admits a strictly positive $\Lambda$-invariant Borel probability measure.

Let $C$ be a unital AH-algebra, let $G$ be a finitely generated abelian group and let $\Lambda : G \to Aut(C)$ be a monomorphism. We show that $C \rtimes_\Lambda G$ can be embedded into a unital simple AF-algebra if and only if $C$ admits a faithful $\Lambda$-invariant tracial state.

## 1 Introduction

Quasidiagonality for crossed product $C^*$-algebras were studied by Dan Voiculescu ([63]), [64] and [65]). Quasidiagonality in $C^*$-algebras has been studied for a long time in many different point of views (see [10] for more information, also [17], [58], [26], [56], [57], [66], [5], [6], [7], [18], [13] and [14], to name a few). Let $X$ be a compact metric space and $\alpha$ be a homeomorphism on $X$. It was proved by Pimsner ([53]) that $C(X) \rtimes_\alpha \mathbb{Z}$ is quasidiagonal if and only if $\alpha$ is pseudo-non-wondering, and if and only if $C(X) \rtimes_\alpha \mathbb{Z}$ can be embedded into an AF-algebra. A more general question is when $A \rtimes_\alpha \mathbb{Z}$ can be embedded into an AF-algebra, where $\alpha \in Aut(A)$ is an automorphism on $A$ and where $A$ is a unital separable $C^*$-algebra which can be embedded into a unital simple AF-algebra. Let $\Lambda$ be a $\mathbb{Z}^2$ action on $X$. Dan Voiculescu (4.6 of [65] ) asked when $C(X) \rtimes_\Lambda \mathbb{Z}^2$ can be embedded into an AF-algebra.

For the first question, there are some significant progresses around the turn of the century. Let $A$ be an AF-algebra and $\alpha \in Aut(A)$ be an automorphism. Nate Brown ([8]) proved (1998) that $A \rtimes_\alpha \mathbb{Z}$ can be embedded into an AF-algebra if and only if $A \rtimes_\alpha \mathbb{Z}$ is quasidiagonal. More importantly, a $K$-theoretical necessary and sufficient condition for $A \rtimes_\alpha \mathbb{Z}$ being quasidiagonal is also given there. When $A$ is a unital simple $A\mathbb{T}$-algebra of real rank zero, Matui (around 2001) ([51]) showed that $A \rtimes_\alpha \mathbb{Z}$ can always be embedded into a unital simple AF-algebra. We find that it is particularly important to know when a crossed product can be embedded into a unital simple AF-algebra. More recently, it was proved ([49]) that, for any unital AH-algebra $A$, $A \rtimes_\alpha \mathbb{Z}$ can be embedded into a unital simple AF-algebra if (and only if) $A$ has a faithful $\alpha$-invariant tracial state. Note that $A$ is not assumed to have real rank zero, nor simple.

It is much more difficult to answer the Voiculescu question above. However, there were some progresses made in last 15 years or so. H. Matui ([51]) showed that under some additional condition on the action, $C(X) \rtimes_\Lambda \mathbb{Z}^2$ can be embedded into an AF-algebra. N. Brown ([9]) proved that if $A$ is a UHF-algebra and $\Lambda : \mathbb{Z}^k \to Aut(A)$ is a homomorphism then $A \rtimes_\Lambda \mathbb{Z}^k$ can be embedded into an AF-algebra.

Suppose that there is a unital monomorphism $h : C(X) \rtimes_\Lambda \mathbb{Z}^k \to A$ for some unital simple AF-algebra. Suppose that $\tau$ is a tracial state on $A$. Then $\tau \circ h$ gives a strictly positive $\Lambda$-invariant probability Borel measure (every non-empty open set has a positive measure). In other words, $\tau \circ h$ is a faithful $\Lambda$-invariant tracial state. The main result of this paper is to show the following theorem: Let $C$ be a unital AH-algebra, let $G$ be a finitely generated abelian group and let $\Lambda : G \to Aut(C)$ be a homomorphism. Then the crossed product $C^*$-algebra $C \rtimes_\Lambda G$ can be embedded into a unital simple AF-algebra if and only if $C$ has a faithful $\Lambda$-invariant tracial state. In particular, when $A = C(X)$ and $G = \mathbb{Z}^k$, this result implies that $C(X) \rtimes_\Lambda \mathbb{Z}^k$ can be embedded into a unital simple AF-algebra if and only if $X$ admits a strictly positive $\Lambda$-invariant Borel probability measure.

In [8], [51], as well as in [10], a version of so-called Berg's technique combined together with a version of Rokhlin property played an important role. Let $C$ be a unital separable amenable $C^*$-algebra and let $\alpha \in Aut(C)$ be an automorphism. Suppose that one can formulate a right condition for two unital monomorphisms $\varphi_1, \varphi_2 : C \to A$, where $A$ is a unital simple AF-algebra, to be asymptotically unitarily equivalent, i.e., there is a continuous path of unitaries $\{u_t : t \in [0, \infty)\}$ of $A$ such that

$$\lim_{t\to\infty} \mathrm{ad}\, u_t \circ \varphi_1(a) = \varphi_2(a)$$

for all $a \in C$. Then the technique developed in the above mentioned articles can be used to prove that $C \rtimes_\alpha \mathbb{Z}$ can be embedded into a unital simple AF-algebra (see 5.5 of [48]). In [49], we present a necessary and a sufficient $K$-theoretical condition for $\varphi_1$ and $\varphi_2$ being asymptotically unitarily equivalent in the case that $C$ is a unital AH-algebra. From this, we obtain a necessary and sufficient condition for $C \rtimes_\alpha \mathbb{Z}$ to be embedded into a unital simple AF-algebra in the case that $C$ is a unital AH-algebra mentioned above (see 10.5 of [49]). However, it is not clear how to formulate a higher dimensional version of Berg's technique.

On the other hand, one may write $C \rtimes_\Lambda \mathbb{Z}^2 = (C \rtimes_{\alpha_1} \mathbb{Z}) \rtimes_{\alpha_2} \mathbb{Z}$ for some automorphisms $\alpha_1$ and $\alpha_2$ on $C$. To prove a true asymptotic unitary equivalence theorem for $C^*$-algebras with the form $C \rtimes_{\alpha_1} \mathbb{Z}$ as above seems to require much more general theory for the classification of amenable $C^*$-algebras. However, we try to pass it via a special path. Given two monomorphisms $\varphi_1, \varphi_2 : C \rtimes_{\alpha_1} \mathbb{Z} \to A$, where $A$ is a unital simple AF-algebra with a unique tracial state and $K_0(A) = \rho_A(K_0(A))$, i.e., $K_0(A)$ has no infinitesimal elements, we will give a (necessary and sufficient) condition for their induced maps $\varphi_1^{(1)}, \varphi_2^{(1)}$ from $C \rtimes_{\alpha_1} \mathbb{Z}$ to $B$, a larger simple AF-algebra, to be asymptotically unitarily equivalent. This will lead to a monomorphism from $C \rtimes_\Lambda \mathbb{Z}^2$ into another unital simple AF-algebra.

To establish such a result, as in the case of AH-algebra (see [49]), it requires some version of so-called Basic Homotopy Lemma for the crossed products. It in turn requires an approximate unitary equivalence theorem for the crossed products. Our strategy is first to prove such a result. This requires a non-commutative version of Berg's technique and a version of Rokhlin property as well as a uniqueness theorem for monomorphisms from an AH-algebra. We also need an existence type theorem to provide the needed bott map. Combining these, we are able to establish the required homotopy lemma. Finally, to obtain similar results beyond $\mathbb{Z}^2$ action, one uses the induction. Several arguments will be repeated.

The paper is organized as follows: Section 2 provides a number of conventions, facts and reviews a couple of known results which will be used several times in the proof. Section 3 is devoted to the approximate unitary equivalence for monomorphisms from the crossed products. In section 4, we show that the prescribed bott map can be constructed using some existing knowledge. Section 5 establishes the required homotopy lemma. In section 6, we proved the needed asymptotic unitary equivalence theorem for the crossed products. In section 7, we present the theorem that $C \rtimes_\Lambda \mathbb{Z}^2$ can be embedded into a unital simple AF-algebra if and only if $C$ admits a faithful $\Lambda$-invariant tracial state. Section 8 provides some absorption lemma. Finally, in section 9, we use the induction to prove the main embedding theorem.

# 2 Preliminaries

**2.1.** Let $A$ be a $C^*$-algebra. Denote by $T(A)$ the tracial state space of $A$ and denote by $Aff(T(A))$ the space of all real affine continuous functions on $T(A)$. Denote by $\rho_A : K_0(A) \to Aff(T(A))$ the positive homomorphism defined by $\rho_A([p]) = \tau \otimes Tr(p)$ for all $\tau \in T(A)$ and projections $p \in M_k(A)$, where $Tr$ is the (non-normalized) standard trace.

**2.2.** Let $U(A)$ be a unitary group of $A$. Denote by $Aut(A)$ the automorphism group. If $u \in U(A)$, denote by $\mathrm{ad}\, u$ the inner automorphism defined by $\mathrm{ad}\, u(a) = u^*au$ for all $a \in A$.

**2.3.** A $C^*$-algebra $A$ is an AH-algebra if $A = \lim_{n\to\infty}(A_n, \psi_n)$, where each $A_n$ has the form $P_nM_{k(n)}(C(X_n))P_n$, where $X_n$ is a finite CW complex (not necessarily connected) and $P_n \in M_{k(n)}(C(X_n))$ is a projection.

We use $\psi_{n,\infty} : A_n \to A$ for the induced homomorphism. Note that every separable commutative $C^*$-algebra is an AH-algebra. AF-algebras and $A\mathbb{T}$-algebras are AH-algebras.

**2.4.** Denote by $\mathcal{N}$ the class of separable amenable $C^*$-algebras which satisfies the Universal Coefficient Theorem ([59]).

**2.5.** Let $X$ be a compact metric space. Denote by $Homeo(X)$ the group of all homeomorphisms on $X$.

**2.6.** Let $A$ and $B$ be two $C^*$-algebras and let $L_1, L_2 : A \to B$ be two maps. Let $\epsilon > 0$ and $\mathcal{F} \subset A$ be a subset. We write

$$L_1 \approx_\epsilon L_2 \text{ on } \mathcal{F},$$

if

$$\|L_1(a) - L_2(a)\| < \epsilon \text{ for all } a \in \mathcal{F}.$$

We say map $L_1$ is $\epsilon$-$\mathcal{F}$-multiplicative if

$$\|L_1(ab) - L_1(a)L_1(b)\| < \epsilon \text{ for all } a, b \in \mathcal{F}.$$

**Definition 2.7.** Let $A$ and $C$ be two unital $C^*$-algebras and let $\varphi_1, \varphi_2 : C \to A$ be two unital monomorphisms. Define the mapping torus

$$M_{\varphi_1,\varphi_2} = \{f \in C([0,1], A) : f(0) = \varphi_1(c), \;\; f(1) = \varphi_2(c) \text{ for some } c \in C\}.$$

One obtains an exact sequence:

$$0 \to SA \xrightarrow{\imath} M_{\varphi,\psi} \xrightarrow{\pi_0} C \to 0. \tag{e 2.1}$$

Denote by $\pi_t : M_{\varphi_1,\varphi_2} \to C$ the point-evaluation at $t$ ($t \in [0,1]$).

Suppose that $C$ is a separable amenable $C^*$-algebra. From (e 2.1), one obtains an element in $Ext(C, SA)$. In this case we identify $Ext(C, SA)$ with $KK^1(C, SA)$ and $KK(C, A)$.

Suppose that $[\varphi] = [\psi]$ in $KK(C, A)$ and $C$ satisfies the Universal Coefficient Theorem, using Dadarlat-Loring's notation, one has the following splitting exact sequence:

$$0 \to \underline{K}(SA) \xrightarrow{[\imath]} \underline{K}(M_{\varphi_1,\varphi_2}) \overset{[\pi_0]}{\rightleftarrows}_{\theta} \underline{K}(C) \to 0. \tag{e 2.2}$$

In other words there is $\theta \in Hom_\Lambda(\underline{K}(C), \underline{K}(M_{\varphi_1,\varphi_2}))$ such that $[\pi_0] \circ \theta = [\mathrm{id_C}]$. In particular, one has a monomorphism $\theta|_{K_1(C)} : K_1(A) \to K_1(M_{\varphi,\psi})$ such that $[\pi_0] \circ \theta|_{K_1(A)} = (\mathrm{id}_C)_{*1}$. Thus, one may write

$$K_1(M_{\varphi_1,\varphi_2}) = K_0(A) \oplus K_1(C). \tag{e 2.3}$$

**Definition 2.8.** Suppose that $T(A) \neq \emptyset$. Let $u \in M_l(M_{\varphi,\psi})$ be a unitary which is a piecewise smooth function on $[0,1]$. For each $\tau \in T(A)$, we denote by $\tau$ for the trace $\tau \otimes Tr$ on $M_l(A)$, where $Tr$ is the standard trace on $M_l$. Define

$$R_{\varphi,\psi}(u)(\tau) = \frac{1}{2\pi i}\int_0^1 \tau(\frac{du(t)}{dt}u(t)^*)dt. \tag{e 2.4}$$

$R_{\varphi,\psi}(u)(\tau)$ is real for every $\tau \in T(A)$.

Suppose that $[\varphi_1] = [\varphi_2]$ in $KL(C, A)$. We also assume that

$$\tau(\varphi_1(c)) = \tau(\varphi_2(c)) \text{ for all } c \in C \text{ and } \tau \in T(A). \tag{e 2.5}$$

Exactly as in section 2 of [30], one has the following statement:

*When* $[\varphi_1] = [\varphi_2]$ *in* $KL(C, A)$ *and* (e 2.5) *holds, there exists a homomorphism*

$$R_{\varphi,\psi} : K_1(M_{\varphi,\psi}) \to Aff(T(A))$$

*defined by*

$$R_{\varphi,\psi}([u])(\tau) = \frac{1}{2\pi i}\int_0^1 \tau(\frac{du(t)}{dt}u(t)^*)dt.$$

We will call $R_{\varphi,\psi}$ the *rotation map* for the pair $\varphi$ and $\psi$.

Suppose also that $\tau \circ \varphi_1 = \tau \circ \varphi_2$ for all $\tau \in T(A)$. Then one obtains the homomorphism

$$R_{\varphi,\psi} \circ \theta|_{K_1(C)} : K_1(C) \to Aff(T(A)). \tag{e 2.6}$$

To keep the same notation as in [30], we write

$$\tilde{\eta}_{\varphi_1,\varphi_2} = 0,$$

if $R_{\varphi,\psi} \circ \theta = 0$, i.e., $\theta(K_1(C)) \in \ker R_{\varphi_1,\varphi_2}$ for some such $\theta$. Thus, $\theta$ also gives the following:

$$\ker R_{\varphi,\psi} = \ker \rho_A \oplus K_1(C).$$

Under the assumption that $[\varphi_1] = [\varphi_2]$ in $KK(C, A)$ and $\tau \circ \varphi_1 = \tau \circ \varphi_2$ for all $\tau \in T(A)$, and if, in addition, $K_i(C)$ is torsion free, such $\theta$ exists whenever $\rho_A(K_0(A)) = R_{\varphi_1,\varphi_2}(K_1(M_{\varphi_1,\varphi_2}))$ and the following splits:

$$0 \to \ker \rho_A \to \ker R_{\varphi_1,\varphi_2} \to K_1(A) \to 0.$$

For further information about rotation maps, see section 2 of [30] and section 3 of [49].

**2.9.** Let $A$ be a unital amenable $C^*$-algebra, let $B$ be a unital $C^*$-algebra and let $h : A \to B$ be a homomorphism. Suppose that $v \in U(B)$. We will refer to [47] for the definition of $\mathrm{Bott}(h, v)$ and $\mathrm{bott}_1(h, v)$.

Given a finite subset $\mathcal{P} \subset \underline{K}(A)$, there exists a finite subset $\mathcal{F} \subset A$ and $\delta_0 > 0$ such that

$$\mathrm{Bott}(h, v)|_{\mathcal{P}}$$

is well defined, if

$$\|[h(a),\, v]\| = \|h(a)v - vh(a)\| < \delta_0 \text{ for all } a \in \mathcal{F}$$

(see 2.10 of [47]).

There is $\delta_1 > 0$ ([50]) such that $\mathrm{bott}_1(u, v)$ is well defined for any pair of unitaries $u$ and $v$ such that $\|[u,\, v]\| < \delta_1$. As in 2.2 of [22], if $v_1, v_2, ..., v_n$ are unitaries such that

$$\|[u,\, v_j]\| < \delta_1/n, \;\; j = 1, 2, ..., n,$$

then

$$\mathrm{bott}_1(u,\, v_1 v_2 \cdots v_n) = \sum_{j=1}^{n} \mathrm{bott}_1(u,\, v_j).$$

We will also use $\boldsymbol{\beta}$ for the usual homomorphism from $K_1(A)$ to $K_0(A \otimes C(\mathbb{T}))$.

See section 2 of [47] for the further information.

**Definition 2.10.** Denote by $\mathcal{U}$ throughout this paper the universal UHF-algebra $\mathcal{U} = \otimes_{n \geq 1} M_n$.

Let $\{e_{i,j}^{(n)}\}$ be the canonical matrix units for $M_n$. Let $u_n \in M_n$ be the unitary matrix such that $\mathrm{ad}\, u_n(e_{i,i}^{(n)}) = e_{i+1,i+1}^{(n)}$ (modulo $n$). Let $\sigma = \otimes_{n \geq 1} \mathrm{ad}\, u_n \in Aut(\mathcal{U})$ be the shift (see for example Example 2.2 of [8]). A fact that we will use in this paper is the following *cyclic Rokhlin property* that $\sigma$ has: For any integer $k > 0$, any $\epsilon > 0$ and any finite subset $\mathcal{F} \subset \mathcal{U}$, there exist mutually orthogonal projections $e_1, e_2, ..., e_k \in \mathcal{U}$ such that $\sum_{i=1}^{k} e_i = 1_{\mathcal{U}}$, $\|x e_i - e_i x\| < \epsilon$ for all $x \in \mathcal{F}$ and $\sigma(e_i) = e_{i+1}, i = 1, 2, ..., k$ $(e_{k+1} = e_1)$.

Denote $\mathcal{U}^k = \overbrace{\mathcal{U} \otimes \mathcal{U} \otimes \cdots \otimes \mathcal{U}}^{k}$. We note that $\mathcal{U}^k \cong \mathcal{U}$.

Throughout this paper, $\mathcal{U}$ and $\sigma$ will be as the above.

**Definition 2.11.** It follows from [51] (and its proof) that there is a unital monomorphism $j : \mathcal{U} \times_\sigma \mathbb{Z} \to \mathcal{U}$. However, in this case, more is true. First $[\sigma] = [\mathrm{id}|_U]$ in $KK(\mathcal{U},\mathcal{U})$ and $\tau = \tau \circ \sigma$. Since $K_1(\mathcal{U}) = \{0\}$, $K_1(M_{\mathrm{id}_{\mathcal{U}},\sigma}) = K_0(\mathcal{U})$. In particular (see [30]), there exists a continuous path of unitaries $\{v(t) : t \in [0, \infty)\}$ of $\mathcal{U}$ such that

$$\lim_{t \to \infty} v(t)^* a v(t) = \sigma(a) \;\text{ for all }\; a \in \mathcal{U}. \tag{e 2.7}$$

Therefore, as in [51], there is a unital embedding $\varphi : \mathcal{U} \rtimes_\sigma \mathbb{Z} \to \mathcal{U}$ such that

$$\tau \circ \varphi = \tau. \tag{e 2.8}$$

Define $\psi : \mathcal{U} \rtimes_\sigma \mathbb{Z} \to \mathcal{U} \otimes \mathcal{U}$ by $\psi(a) = \varphi(a) \otimes 1_{\mathcal{U}}$ for all $a \in \mathcal{U}$ and $\psi(u_\sigma) = \varphi(u_\sigma) \otimes \varphi(u_\sigma^*)$. Then $\psi$ is a unital monomorphism. Denote by $s : \mathcal{U} \otimes \mathcal{U} \to \mathcal{U}$ an isomorphism with $s_{*0} = \mathrm{id}_{K_0(\mathcal{U} \otimes \mathcal{U})}$. We define $\imath : \mathcal{U} \rtimes_\sigma \mathbb{Z} \to \mathcal{U}$ by $s \circ \psi$.

*In what follows, $\imath$ will be used without further explanation.*

**Lemma 2.12.** *Let $C$ be a unital separable $C^*$-algebra and let $\alpha \in Aut(C)$ be an automorphism. Let $B$ be a unital separable amenable $C^*$-algebra. Suppose that $\varphi : C \rtimes_\alpha \mathbb{Z} \to B$ is a unital homomorphism such that $\varphi|_C$ is injective. Define a unital homomorphism $\varphi^{(1)} : C \rtimes_\alpha \mathbb{Z} \to B \otimes \mathcal{U}$ defined by $\varphi^{(1)}(c) = \varphi(c) \otimes 1_{\mathcal{U}}$ for all $c \in C$ and $\varphi^{(1)}(u_\alpha) = \varphi(u_\alpha) \otimes \imath(u_\sigma)$.*

*Then $\varphi^{(1)}$ is a monomorphism.*

*Proof.* Let $C_1 = (C \otimes \mathcal{U}) \rtimes_{\alpha \otimes \sigma} \mathbb{Z}$. Then $\alpha \otimes \sigma$ is known to have the cyclic Rokhlin property (see 2.10). We view $\mathcal{U}$ as a $C^*$-subalgebra of $\mathcal{U} \rtimes_\sigma \mathbb{Z}$. Define $\psi : C_1 \to B \otimes \mathcal{U}$ by $\psi(c \otimes b) = \varphi(c) \otimes \imath(b)$ for all $c \in C$ and $b \in \mathcal{U}$ and $\psi(u_{\alpha \otimes \sigma}) = \varphi(u_\alpha) \otimes \imath(u_\sigma)$. Then $C_1|_{C \otimes \mathcal{U}}$ is injective. Since $\alpha \otimes \sigma$ has the cyclic Rokhlin property, by Lemma 4.1 of [46], $\psi$ is injective. Let $j_0 : C \rtimes_\alpha \mathbb{Z} \to C_1$ be defined by $j_0(c) = c \otimes 1_{\mathcal{U}}$ for $c \in C$ and $j_0(u_\alpha) = u_{\alpha \otimes \sigma}$. Then $j_0$ is an embedding. Note $\varphi^{(1)} = \psi \circ j_0$.

□

**Theorem 2.13.** (Corollary 4.8 of [44]) *Let $C$ be a unital AH-algebra and let $\tau_0 \in T(C)$ be a faithful tracial state.*

*Let $\epsilon > 0$, $\mathcal{F} \subset C$ be a finite subset. There exists $\delta > 0$, $\sigma > 0$, a finite subset $\mathcal{G}$ and a finite subset $\mathcal{P} \subset \underline{K}(C)$ satisfying the following:*

*For any unital simple $C^*$-algebra with tracial rank zero with a unique tracial state $\tau \in T(A)$, any two unital $\delta$-$\mathcal{G}$-multiplicative contractive completely positive linear maps $L_1, L_2 : C \to A$ such that*

$$[L_1]|_{\mathcal{P}} = [L_2]|_{\mathcal{P}} \text{ and} \tag{e 2.9}$$

$$\tau \circ L_i \approx_{\sigma} \tau_0 \ \textit{on} \ \mathcal{G}, \tag{e 2.10}$$

*there exists a unitary $w \in U(A)$ such that*

$$\mathrm{ad}\, w \circ L_1 \approx_{\epsilon} L_2 \ \textit{on} \ \mathcal{F}. \tag{e 2.11}$$

*Proof.* It is clear that the general case can be reduced to the case that $C = PM_k(C(X))P$, where $X$ is a finite-dimensional compact metric space and $P \in M_k(C(X))$ is a projection. Since $\tau_0$ is fixed, the statement follows from Cor. 4.8 of [44].

□

The following is a direct application of a theorem Choi and Effros ([12], see 5.10.10 of [35]).

**Proposition 2.14.** *Let $A$ be a separable amenable $C^*$-algebra. For any $\epsilon > 0$ and finite subset $\mathcal{F} \subset A$, there is $\delta > 0$ and a finite subset $\mathcal{G} \subset A$ satisfying the following:*

*For any $C^*$-algebra $B$ and any self-adjoint linear map $L : A \to B$ which is $\delta$-$\mathcal{G}$-multiplicative, there exists an $\epsilon$-$\mathcal{F}$-multiplicative contractive completely positive linear map $\varphi : A \to B$ such that*

$$L \approx_{\epsilon} \varphi \ \textit{on} \ \mathcal{F}. \tag{e 2.12}$$

*If, furthermore, $A$ has a unital, one may assume that $\varphi(1_A)$ is a projection.*

## 3 Approximate unitary equivalence

**Lemma 3.1.** *Let $C$ be a unital amenable separable $C^*$-algebra and $\alpha \in Aut(C)$ be an automorphism. For any $\epsilon > 0$, $L > 0$ and finite subset $\mathcal{F} \subset C$, there is $\delta > 0$ and a finite subset $\mathcal{G} \subset C$ satisfying the following:*

*Suppose that $A$ and $B$ are two unital amenable separable $C^*$-algebras, $\varphi : C \rtimes_{\alpha} \mathbb{Z} \to A$ and $\psi : \mathcal{U} \rtimes_{\sigma} \mathbb{Z} \to B$ are two unital monomorphisms. Fix unitaries $u_1 \in A$, $u_2 \in B$ such that $u_1^* \varphi(a) u_1 = \varphi(\alpha(a))$ for all $a \in C$ and $u_2^* \psi(b) u_2 = \psi(\sigma(b))$ for all $b \in \mathcal{U}$. Suppose that there is a unitary $v \in U(A)$ and a continuous path of unitaries $\{v_t : t \in [0,1]\} \subset A \otimes 1$ such that*

$$v_0 = 1, \ \ v_1 = \prod_{j=0}^{n-1} (u^{n-1-j})^* (v \otimes 1_B) u^{n-1-j}, \tag{e 3.13}$$

$$\|[v,\, \varphi(a)]\| < \delta, \ \ \|[v_t,\, \varphi(a) \otimes 1_B]\| < \delta \ \textit{ for all } \ a \in \mathcal{G} \ \textit{ and } \ t \in [0,1], \tag{e 3.14}$$

$$\textit{and } Length(\{u_t\}) \le L \ \text{ and } \ (L+1)/n < \epsilon/2, \tag{e 3.15}$$

*where $u = u_1 \otimes u_2$, then there is a unitary $w \in A \otimes B$ such that*

$$\mathrm{ad}\, w \circ \varphi(a) \approx_{\epsilon} \varphi(a) \ \textit{ for all } \ a \in \mathcal{F} \ \text{ and } \ w^* u (v \otimes 1_B) w \approx_{\epsilon} u, \tag{e 3.16}$$

*where we identify $a$ with $a \otimes 1$ for $a \in A$.*

*Proof.* Fix $\epsilon > 0$. We may assume that $\mathcal{F}$ is in the unit ball of $C$. Put $\delta = \frac{\epsilon}{4n(n+1)}$. Let $\mathcal{G} = \cup_{-2n\le j\le 2n}\alpha^j(\mathcal{F})$. Suppose that $\{v_t\}$ satisfies the assumption.

There are mutually orthogonal projections $q_1, q_2, ..., q_n \in \psi(\mathcal{U})$ with $\sum_{j=1}^n q_j = 1_{\mathcal{U}}$ such that

$$u_2^* q_j u_2 = q_{j+1}, \;\; j = 1, 2, ...., n \qquad (q_{n+1} = q_1). \tag{e 3.17}$$

Define $e_i = 1 \otimes q_i$, $i = 1, 2, ..., n+1$ with $e_{n+1} = e_1$.

In what follows, we will identify an element $a \in A$ with the element $a \otimes 1$ in $A \otimes B$ whenever it is convenient. So we may write that $v = v \otimes 1$ and note that

$$v_1 = ((u^{n-1})^* v u^{n-1})((u^{n-2})^* v u^{n-2})((u^{n-3})^* v(u^{n-3})\cdots u^* v u v.$$

There are $y_1, y_2, ..., y_n \in \{v_t\} \subset A \otimes 1$ such that

$$y_1 = 1, \;\; y_n = v_1 \;\text{ and }\; \|y_j - y_{j+1}\| < \frac{L}{n}, \;\; j = 1, 2, ..., n. \tag{e 3.18}$$

Put $x_1 = y_1^*$, $x_2 = y_1 y_2^*$, $x_3 = y_2 y_3^*$,..., $x_{n-1} = y_{n-2} y_{n-1}^*$ and $x_n = y_{n-1} y_n^*$. Note that

$$\|[\varphi(a),\, x_j]\| < 2\delta \;\text{ for all }\; a \in \mathcal{G}, \;\; j = 1, 2, ..., n. \tag{e 3.19}$$

Put

$$s_j = v^* u^* v u \cdots (u^{j-1})^* v^* u^{j-1}, \;\; j = 1, 2, ..., n-1$$

Define

$$\eta_1 = u^{n-1} s_{n-1}^* x_1 s_{n-1} (u^{n-1})^* \tag{e 3.20}$$
$$\eta_2 = u^{n-2} s_{n-2}^* x_2 s_{n-2} (u^{n-2})^*, \tag{e 3.21}$$
$$\cdots, \tag{e 3.22}$$
$$\eta_{n-2} = u^2 s_2^* x_{n-2} s_2 (u^2)^* = u^2 (u^* v u) v x_{n-2} v^* (u^* v^* u)(u^2)^* \tag{e 3.23}$$
$$\eta_{n-1} = u v x_{n-1} v^* u^* \tag{e 3.24}$$
$$\eta_n = x_n. \tag{e 3.25}$$

Note that

$$\varphi(a) u v u^* = u u^* \varphi(a) u v u^* = u \varphi\circ\alpha(a) v u^* \tag{e 3.26}$$
$$\approx_\delta u v \varphi\circ\alpha(a) u^* \tag{e 3.27}$$
$$= u v u^* u \varphi\circ\alpha(a) u^* = u v u^* \varphi(a) \tag{e 3.28}$$

for all $a \in \cup_{-2(n-1)\le j\le 2(n-1)}\alpha(\mathcal{F})$. It follows that

$$\|[\varphi(a),\, \eta_j]\| < 2(n-j+1)\delta \;\text{ for all }\; a \in \cup_{-2(n-j)\le j\le 2(n-j)}\alpha^j(\mathcal{F}) \tag{e 3.29}$$

and for $j = 1, 2, ..., n$. Put

$$d_1 = e_1 v \eta_1 e_1, \;\; d_2 = e_2 v \eta_2 e_2, \;\; d_3 = u^* d_2 u v \eta_3 e_3, ... \tag{e 3.30}$$
$$d_{j+1} = u^* d_j u v \eta_{j+1} e_{j+1}, ..., d_n = u^* d_{n-1} u v \eta_n e_n. \tag{e 3.31}$$

Note that

$$d_n = (u^2)^* d_{n-2} u^2 u^* v (u v x_{n-1} v^* u^*) u v \eta_n e_n \tag{e 3.32}$$
$$= (u^2)^* d_{n-2} u^2 u^* v u v x_{n-1} x_n e_n \tag{e 3.33}$$
$$= (u^3)^* d_{n-3} u^3 (u^2)^* v (u^2 u^* v u v x_{n-2} v^* (u^* v u)(u^2)^*)(u^2 u^* v u v x_{n-1} x_n e_n) \tag{e 3.34}$$
$$= (u^3)^* d_{n-3} u^3 (u^2)^* v u^2 u^* v u v x_{n-2} x_{n-1} x_n e_n \tag{e 3.35}$$
$$\cdots \tag{e 3.36}$$
$$= v_1 x_1 x_2 \cdots x_n e_n = e_n \tag{e 3.37}$$

Define

$$Z=\sum_{j=1}^{n} d_j. \tag{e 3.38}$$

Note that $Z\in U(A\otimes B)$. We estimate that

$$\varphi(a)d_1 \quad = \quad e_1\varphi(a)ve_1\eta_1e_1 \tag{e 3.39}$$
$$\approx_{\delta} \quad e_1v\varphi(a)\eta_1e_1 \tag{e 3.40}$$
$$\approx_{2n\delta} \quad e_1v\eta_1e_1\varphi(a)=d_1\varphi(a) \tag{e 3.41}$$

for all $a\in\mathcal{G}$. Suppose that we have shown

$$\varphi(a)d_j \approx_{(j(2n+3)-j(j+1))\delta} d_j\varphi(a) \ \text{ for all } \ a\in\cup_{-2(n-j)\le i\le 2(n-j)}\alpha^i(\mathcal{F}). \tag{e 3.42}$$

$$\varphi(a)d_{j+1} \quad = \quad \varphi(a)u^*d_juv\eta_je_{j+1} \tag{e 3.43}$$
$$= \quad u^*\varphi\circ\alpha^{-1}(a)d_juv\eta_je_{j+1} \tag{e 3.44}$$
$$\approx_{(j(2n+3)-j(j+1))\delta} \quad u^*d_j\varphi\circ\alpha^{-1}(a)uv_1\eta_je_{j+1} \tag{e 3.45}$$
$$= \quad u^*d_ju\varphi(a)v\eta_je_{j+1} \tag{e 3.46}$$
$$\approx_{\delta} \quad u^*d_juv\varphi(a)\eta_je_{j+1} \tag{e 3.47}$$
$$\approx_{2(n-j)\delta} \quad u^*d_juv\eta_je_{j+1}\varphi(a) \tag{e 3.48}$$

for all $a\in\cup_{-2(n-j-1)\le i\le 2(n-j-1)}\alpha^i(\mathcal{F})$. In other words,

$$\varphi(a)d_{j+1}\approx_{(j+1)(2n+3)-(j+1)(j+2)\delta} d_{j+1}\varphi(a) \tag{e 3.49}$$

for all $a\in\cup_{-2(n-j-1)\le i\le 2(n-j-1)}\alpha^i(\mathcal{F})$. It follows that

$$\|[Z,\,\varphi(a)]\|<2n(n+1)\delta \ \text{ for all } \ a\in\mathcal{F}. \tag{e 3.50}$$

Note that

$$\|\eta_j-1\|<L/n<\epsilon/2,\ \ j=1,2,...,n. \tag{e 3.51}$$

We then compute that

$$Z^*uZ \quad = \quad d_n^*e_nue_1d_1+\sum_{j=1}^{n-1}d_j^*e_jue_{j+1}d_{j+1} \tag{e 3.52}$$
$$= \quad e_nuv\eta_1e_1+\sum_{j=1}^{n-1}u(u^*d_j^*ue_{j+1}d_{j+1}) \tag{e 3.53}$$
$$= \quad e_nuv\eta_1e_1+\sum_{j=1}^{n-1}u(u^*d_j^*ue_{j+1}u^*d_juv\eta_je_{j+1}) \tag{e 3.54}$$
$$= \quad e_nuv\eta_1e_1+\sum_{j=1}^{n-1}ue_{j+1}v\eta_je_{j+1} \tag{e 3.55}$$
$$\approx_{\epsilon} \quad e_nuve_1+\sum_{j=1}^{n-1}uve_{j+1}=uv \qquad \text{(applying(e 3.51))}. \tag{e 3.56}$$

Put $w = Z^*$. It follows from (e 3.50) that

$$\operatorname{ad} w \circ \varphi(a) \approx_\epsilon \varphi(a) \ \text{ for all } \ a \in \mathcal{F}. \tag{e 3.57}$$

Moreover, by (e 3.52),

$$w^*(u_1 v \otimes u_2)w = Z(u_1 v \otimes u_2)Z^* \approx_\epsilon u. \tag{e 3.58}$$

$\square$

**Lemma 3.2.** *Let $C$ be a unital AH-algebra and let $\alpha \in Aut(C)$ be an automorphism. Suppose that $A$ and $B$ are two unital simple $C^*$-algebras with tracial rank zero, $\varphi : C \rtimes_\alpha \mathbb{Z} \to A$ and $\psi : \mathcal{U} \rtimes_\sigma \mathbb{Z} \to B$ are two unital monomorphisms.*

*For any $\epsilon > 0$ and any finite subset $\mathcal{F} \subset C$, there is $\delta > 0$, a finite subset $\mathcal{G} \subset C$ and a finite subset $\mathcal{P} \subset \underline{K}(C)$ satisfying the following:*

*Fix unitaries $u_1 \in A$, $u_2 \in B$ such that $u_1^*\varphi(a)u_1 = \varphi(\alpha(a))$ for all $a \in C$ and $u_2^*\psi(b)u_2 = \psi(\sigma(b))$ for all $b \in \mathcal{U}$. Suppose that there is a unitary $v \in U(A)$ such that*

$$\|[v,\, \varphi(a)]\| < \delta, \quad \textit{for all } a \in \mathcal{G} \ \textit{and} \ \mathrm{Bott}(\varphi,\, v)|_{\mathcal{P}} = 0, \tag{e 3.59}$$

*then there is a unitary $w \in A \otimes B$ such that*

$$w^*(\varphi(a) \otimes 1_B)w \approx_\epsilon \varphi(a) \otimes 1_B \ \textit{for all } a \in \mathcal{F} \ \text{and} \ w^*u(v \otimes 1_B)w \approx_\epsilon u, \tag{e 3.60}$$

*where $u = u_1 \otimes u_2$.*

*Proof.* Fix a finite subset $\mathcal{F} \subset C$ and $\epsilon > 0$. Define $\varphi' : C \to A \otimes B$ by $\varphi'(c) = \varphi(c) \otimes 1_B$ for all $c \in C$.

Let $\delta_1 > 0$ (in place of $\delta$) and $\mathcal{G}_1 \subset C$ (in place of $\mathcal{G}$) be a finite subset required by 3.1 for $L = 2\pi + 1$.

Let $n \geq 1$ be an integer such that $\frac{2\pi+1}{n} < \min\{\epsilon/2, \delta_1/2\}$.

Let $\delta_2 > 0$ (in place of $\delta$), $\mathcal{G}_2 \subset C$ (in place of $\mathcal{G}$) be a finite subset and let $\mathcal{P}_1 \subset \underline{K}(C)$ be a finite subset required by 17.9 of [47] corresponding to $\delta_1/2$ (in place of $\epsilon$), $\mathcal{G}_1$ (in place of $\mathcal{F}$) and $\varphi'|_C$.

Without loss of generality, to simplify notation, we may assume that $\mathcal{F}$, $\mathcal{G}_1$ and $\mathcal{G}_2$ are all in the unit ball of $C$. We may further assume that $\delta_2 < \min\{\delta_1/2, \epsilon/2\}$.

Define

$$v_1 = \prod_{j=0}^{n-1} (u^{n-1-j})^*(v \otimes 1)u^{n-1-j}, \tag{e 3.61}$$

where $u = u_1 \otimes u_2$. Note that $v_1 \in A \otimes 1$. Define

$$\mathcal{G} = \cup_{k=-2n}^{2n} \alpha^k(\mathcal{G}_2) \ \text{ and } \ \mathcal{P} = \cup_{k=-2n}^{2n} [\alpha^k](\mathcal{P}_1).$$

Let $\delta > 0$ so that $\delta < \delta_2/2n^2$.

Suppose that

$$\|[\varphi(c),\, v]\| < \delta \ \text{ for all } \ c \in \mathcal{G} \ \text{ and } \ \mathrm{Bott}(\varphi,\, v)|_{\mathcal{P}} = 0. \tag{e 3.62}$$

We may also assume that $\mathrm{Bott}(h', v')|_{\mathcal{P}}$ is well defined provided that

$$\|[h',\, v']\| < n^2\delta < \delta_2/2 \tag{e 3.63}$$

for any unital homomorphism $h'$ from $C$ and any unitary $v'$.

Note we have

$$\mathrm{Bott}(\varphi',\, v\otimes 1_B)|_{\mathcal{P}} = 0. \tag{e 3.64}$$

Note that,

$$\begin{aligned} \varphi'(a)u^*(v\otimes 1_B)u &= u^*\varphi'(\alpha^{-1}(a))(v\otimes 1_B)u && \text{(e 3.65)}\\ &\approx_\delta u^*(v\otimes 1_B)\varphi'(\alpha^{-1}(a))u && \text{(e 3.66)}\\ &= u^*(v\otimes 1_B)u\varphi'(a) && \text{(e 3.67)}\end{aligned}$$

for all $a\in \cup_{k=-2n+1}^{2n-1}\alpha^k(\mathcal{G}_2)$. It follows that

$$\|[\varphi'(a),\, v_1]\| < n\delta < \delta_2/2n \text{ for all } a\in \mathcal{G}_2. \tag{e 3.68}$$

By (e 3.64), we have that

$$\mathrm{Bott}(\varphi',\, (u^j)^*(v\otimes 1_B)u^j)|_{\mathcal{P}_1} = \mathrm{Bott}(\varphi', v\otimes 1_B)|_{[\alpha^{-j}](\mathcal{P}_1)} = 0, \tag{e 3.69}$$

$$\tag{e 3.70}$$

$j = 0, 1, 2, ..., n-1$. It follows that

$$\mathrm{Bott}(\varphi',\, v_1)|_{\mathcal{P}_1} = \sum_{j=1}^{n-1}\mathrm{Bott}(\varphi',\, (u^j)^*(v\otimes 1_B)u^j)|_{\mathcal{P}_1} = 0 \tag{e 3.71}$$

The above is computed in $A\otimes 1$. It follows from the Basic Homotopy Lemma 17.9 of [47] that there exists a continuous path of unitaries $\{v(t) : t\in[0,1]\}\subset U(A\otimes 1)$ such that

$$v(0) = 1,\;\; v(1) = v_1 \text{ and } \|[\varphi'(c),\, v(t)]\| < \delta_1/2 \text{ for all } a\in\mathcal{G}_1. \tag{e 3.72}$$

Moreover

$$\mathrm{Length}(\{v(t)\}) \le 2\pi + 1 = L. \tag{e 3.73}$$

Thus 3.1 applies.

□

**Corollary 3.3.** *Let $C$ be a unital AH-algebra, let $\alpha\in Aut(C)$ be an automorphism and let $A\cong A\otimes\mathcal{U}$ be a unital simple AF-algebra with a unique tracial state $\tau$ and $K_0(A) = \rho_A(K_0(A))$. Suppose that $\varphi_1, \varphi_2 : C\rtimes_\alpha\mathbb{Z}\to A$ are two unital monomorphisms. For any $\epsilon > 0$ and any finite subset $\mathcal{F}\subset C\rtimes_\alpha\mathbb{Z}$, there exists $\delta > 0$, a finite subset $\mathcal{G}\subset C$ and a finite subset $\mathcal{P}\subset K_1(C)$ satisfying the following:*

*Suppose that there exists a unitary $z\in A$ such that*

$$\mathrm{ad}\, z\circ\varphi_1(a)\approx_\delta \varphi_2(a) \text{ for all } a\in\mathcal{G} \tag{e 3.74}$$

*and, for any $x\in\mathcal{P}$,*

$$\mathrm{bott}_1(\varphi_2,\, \varphi_2(u_\alpha)^*z^*\varphi_1(u_\alpha)z)(x) = 0. \tag{e 3.75}$$

*Then there exists a unitary $w\in U(A\otimes\mathcal{U})$ such that*

$$\mathrm{ad}\, w\circ\varphi_1^{(1)} \approx_\epsilon \varphi_2^{(1)} \text{ on } \mathcal{F}, \tag{e 3.76}$$

*where $\varphi_i^{(1)} : C\rtimes_\alpha\mathbb{Z}\to A\otimes\mathcal{U}$ is defined by $\varphi_i^{(1)}(c) = \varphi_i(c)\otimes 1_{\mathcal{U}}$ and $\varphi_i^{(1)}(u_\alpha) = \varphi_i(u_\alpha)\otimes\imath(u_\sigma)$, $i = 1, 2$.*

*Proof.* Let $\epsilon > 0$ and $\mathcal{F} \subset C \rtimes_\alpha \mathbb{Z}$ be a finite subset. Without loss of generality, we may assume that $\mathcal{F} = \mathcal{F}_0 \cup \{u_\alpha\}$, where $\mathcal{F}_0 \subset C$ is a finite subset.

Let $\delta > 0$, $\mathcal{G} \subset C$ be a finite subset and $\mathcal{P}_1 \subset \underline{K}(C)$ be a finite subset required by 3.2 for $\epsilon/2$ and $\mathcal{F}_0$ above (with $\varphi$ being replaced by $\varphi_2$, $B = \mathcal{U}$ and $\psi$ being replaced by $\imath$). Without loss of generality, we may assume that $\mathcal{G}$ is in the unit ball of $C$ and $\delta < \epsilon/2$.

Since $K_0(A)$ is divisible and torsion free, there is a finite subset $\mathcal{P} \subset K_1(C)$ such that

$$\mathrm{Bott}(\varphi_2,\, v)|_{\mathcal{P}_1} = 0 \tag{e 3.77}$$

provided that

$$\mathrm{bott}_1(\varphi_2, v)|_{\mathcal{P}} = 0 \tag{e 3.78}$$

and $v$ is a unitary and $\mathrm{Bott}(\varphi_2,\, v)|_{\mathcal{P}_1}$ is well-defined.

Now suppose that (e 3.74) and (e 3.75) hold for above $\delta$, $\mathcal{G}$ and $\mathcal{P}$.

Define

$$v = \varphi_2(u_\alpha)^* z^* \varphi_1(u_\alpha) z. \tag{e 3.79}$$

Then

$$\|[\varphi_2(c),\, v]\| < \delta \ \text{ for all } \ c \in \mathcal{G} \ \text{ and } \ \mathrm{Bott}(\varphi_2,\, v)|_{\mathcal{P}_1} = 0. \tag{e 3.80}$$

It follows from 3.2 that there exists a unitary $w_1 \in A \otimes \mathcal{U}$ such that

$$\mathrm{ad}\, w_1 \circ \varphi_2^{(1)} \approx_{\epsilon/2} \varphi_2^{(1)} \ \text{ on } \ \mathcal{F}_0 \ \text{ and} \tag{e 3.81}$$

$$w_1^* \varphi_2^{(1)}(u_\alpha)(v \otimes 1_{\mathcal{U}}) w_1 \approx_{\epsilon/2} \varphi_2^{(1)}(u_\alpha). \tag{e 3.82}$$

Note that

$$\varphi_2^{(1)}(u_\alpha)(v \otimes 1_{\mathcal{U}}) = \varphi_2(u_\alpha) v \otimes \imath(u_\sigma) \tag{e 3.83}$$

$$\approx_\delta z^* \varphi_1(u_\alpha) z \otimes \imath(u_\sigma). \tag{e 3.84}$$

Put $w = (z \otimes 1_{\mathcal{U}}) w_1$. Then, by (e 3.81), (e 3.74) and (e 3.83),

$$\mathrm{ad}\, w \circ \varphi_1^{(1)} \approx_\epsilon \varphi_2^{(1)} \ \text{ on } \ \mathcal{F}_0 \ \text{ and } \ \mathrm{ad}\, w \circ \varphi_1^{(1)}(u_\alpha) \approx_\epsilon \varphi_2^{(1)}(u_\alpha). \tag{e 3.85}$$

□

**Corollary 3.4.** *Let $C$ be a unital AH-algebra, let $\alpha \in Aut(C)$ be an automorphism and let $A \cong A \otimes \mathcal{U}$ be a unital simple AF-algebra with a unique tracial state $\tau$ and $K_0(A) = \rho_A(K_0(A))$. Suppose that $\varphi_1, \varphi_2 : C \rtimes_\alpha \mathbb{Z} \to A$ are two unital monomorphisms and suppose that there exists a sequence of unitaries $z_n \in A$ such that*

$$\lim_{n\to\infty} \mathrm{ad}\, z_n \circ \varphi_1(a) = \psi_2(a) \ \text{ for all } \ a \in C \tag{e 3.86}$$

*and, for any $x \in K_1(C)$ and*

$$\mathrm{bott}_1(\varphi_2,\, \varphi_2(u_\alpha)^* z_n^* \varphi_1(u_\alpha) z_n)(x) = 0 \tag{e 3.87}$$

*for all sufficiently large $n$.*

*Then there exists a sequence of unitaries $\{w_n\} \subset U(A \otimes \mathcal{U})$ such that*

$$\lim_{n\to\infty} \mathrm{ad}\, w_n \circ \varphi_1^{(1)}(a) = \varphi_2^{(1)}(a) \ \text{ for all } \ a \in C \rtimes_\alpha \mathbb{Z}, \tag{e 3.88}$$

*where $\varphi_i^{(1)} : C \rtimes_\alpha \mathbb{Z} \to A \otimes \mathcal{U}$ is defined by $\varphi_i^{(1)}(u_\alpha) = \varphi_i(u_\alpha) \otimes \imath(u_\sigma)$, $i = 1, 2$.*

**Theorem 3.5.** *Let $C$ be a unital AH-algebra with $K_1(C)=\{0\}$, let $\alpha\in Aut(C)$ be an automorphism and let $A\cong A\otimes\mathcal{U}$ be a unital simple AF-algebra with a unique tracial state $\tau$ and $K_0(A)=\rho_A(K_0(A))$. Suppose that $\varphi_1,\varphi_2: C\rtimes_\alpha\mathbb{Z}\to A$ are two unital monomorphisms such that*

$$\tau\circ\varphi_1=\tau\circ\varphi_2. \tag{e 3.89}$$

*Then there exists a sequence of unitaries $\{w_n\}\subset U(A\otimes\mathcal{U})$ such that*

$$\lim_{n\to\infty}\operatorname{ad} w_n\circ\varphi_1^{(1)}(a)=\varphi_2^{(1)}(a)\ \ \textit{for all}\ \ a\in C\rtimes_\alpha\mathbb{Z}. \tag{e 3.90}$$

*where $\varphi_i^{(1)}: C\rtimes_\alpha\mathbb{Z}\to A\otimes\mathcal{U}$ by $\varphi_i^{(1)}(c)=\varphi_i(c)\otimes 1_{\mathcal{U}}$ for all $c\in C$ and $\varphi_i^{(1)}(u_\alpha)=\varphi_i(u_\alpha)\otimes\imath(u_\sigma)$, $i=1,2$.*

*Proof.* Since $K_0(A)=\rho_A(K_0(A))$, (e 3.89) implies that

$$(\varphi_1|_C)_{*0}=(\varphi_2|_C)_{*0}. \tag{e 3.91}$$

Since $K_1(A)=\{0\}$ and $K_0(A)$ is divisible, (e 3.91) implies that

$$[\varphi_1|_C]=[\varphi_2|_C]\ \ \text{in}\ \ KK(C,A). \tag{e 3.92}$$

Combining (e 3.92) and (e 3.89), by applying 2.13, there exists a sequence of unitaries $\{z_n\}\in U(A)$ such that

$$\lim_{n\to\infty}\operatorname{ad}z_n\circ\varphi_1(c)=\varphi_2(c)\ \ \text{for all}\ \ c\in C. \tag{e 3.93}$$

Since $K_1(C)=0$,

$$\text{bott}_1(\varphi_2,\,\varphi_2(u_\alpha)^*z_n^*\varphi_1(u_\alpha)z_n)=0. \tag{e 3.94}$$

We then apply 3.4. $\square$

**Theorem 3.6.** *Let $C$ be a unital AH-algebra, let $\alpha\in Aut(C)$ be an automorphism and let $A\cong A\otimes\mathcal{U}$ be a unital simple AF-algebra with a unique tracial state $\tau$ and $K_0(A)=\rho_A(K_0(A))$. Suppose that $\varphi_1,\varphi_2: C\rtimes_\alpha\mathbb{Z}\to A$ are two unital monomorphisms such that*

$$\tau\circ\varphi_1=\tau\circ\varphi_2. \tag{e 3.95}$$

*Suppose also that*

$$R_{\varphi_1\circ j_0,\varphi_2\circ j_0}(K_1(C))\subset\rho_A(K_0(A)), \tag{e 3.96}$$

*where $j_0: C\to C\rtimes_\alpha\mathbb{Z}$ is the embedding.*

*Then there exists a sequence of unitaries $\{w_n\}\subset U(A\otimes\mathcal{U})$ such that*

$$\lim_{n\to\infty}\operatorname{ad} w_n\circ\varphi_1^{(1)}(a)=\varphi_2^{(1)}(a)\ \ \textit{for all}\ \ a\in C\rtimes_\alpha\mathbb{Z}. \tag{e 3.97}$$

*where $\varphi_i^{(1)}: C\rtimes_\alpha\mathbb{Z}\to A\otimes\mathcal{U}$ by $\varphi_i^{(1)}(c)=\varphi_i(c)\otimes 1$ for all $c\in C$ and $\varphi_i^{(1)}(u_\alpha)=\varphi_i(u_\alpha)\otimes\imath(u_\sigma)$, $i=1,2$. Then*

*Proof.* Exactly as in the proof of 3.5, we have

$$[\varphi_1|_C] = [\varphi_2|_C] \text{ in } KK(C, A). \tag{e 3.98}$$

Define

$$M_{\varphi_1,\varphi_2} = \{f \in C([0,1], A) : f(0) = \varphi_1(a) \text{ and } f(1) = \varphi_2(a) \text{ for some } a \in C\}.$$

Then the condition (e 3.96) implies that

$$R_{\varphi_1,\varphi_2}(K_1(M_{\varphi_1,\varphi_2})) = \rho_A(K_0(A)). \tag{e 3.99}$$

Then, by combining (e 3.92), by applying Lemma 11.3 of [49] and Theorem 10.7 of [49], one obtains a continuous path of unitaries $\{z_t : t \in [0,\infty)\} \subset U(A)$ such that

$$\lim_{t\to\infty} \operatorname{ad} z_t \circ \varphi_1(a) = \varphi_2(a) \text{ for all } a \in C. \tag{e 3.100}$$

As in the proof of 3.5, put

$$v_t = \varphi_2(u_\alpha) z_t^* \varphi_1(u_\alpha) z_t \text{ for all } t \in [0,\infty). \tag{e 3.101}$$

Then one obtains

$$\lim_{t\to\infty} \|[\varphi_2(a),\, v_t]\| = 0 \text{ for all } a \in C. \tag{e 3.102}$$

In particular, for a fixed finite subset $\mathcal{P} \subset K_1(C)$,

$$\text{bott}_1(\varphi_2,\, v_t)|_{\mathcal{P}} = \text{bott}_1(\varphi_2,\, v_{t'})|_{\mathcal{P}} \tag{e 3.103}$$

for all sufficiently large $t, t'$. By the Exel formula (3.6 of [49]), for each $z \in U(M_k(C))$, one has

$$\lim_{t\to\infty} \text{bott}_1(\varphi_2, v_t)([z])(\tau) = \lim_{t\to\infty} \tau(\log(V_t^* \varphi(z) V_t \varphi(z)^*) = 0, \tag{e 3.104}$$

where $V_t = \text{diag}(\overbrace{v_t, v_t, ..., v_t}^{k})$. Combining (e 3.103) and (e 3.104), one must have, for any $x \in \mathcal{P}$,

$$\text{bott}_1(\varphi_2, v_1)(x)(\tau) = 0. \tag{e 3.105}$$

Since $K_0(A) = \rho_A(K_0(A))$, this implies that

$$\text{bott}_1(\varphi_2, v_t)(x) = 0 \tag{e 3.106}$$

for all $x \in \mathcal{P}$ for all large $t$. Then 3.4 applies.

□

# 4 Bott maps

**Theorem 4.1.** *Let $C$ be a unital separable amenable $C^*$-algebra in $\mathcal{N}$ which has a unital embedding $j : C \to A$ for some unital simple AF-algebra $A$.*

*Then there is a unital separable simple $C^*$-algebra $B \in \mathcal{N}$ with tracial rank zero and a unital monomorphism $\varphi : C \to B$ such that $\varphi_{*1} = \gamma \circ \kappa$, where $\kappa : K_1(C) \to K_1(C)/\mathrm{Tor}(\mathrm{K_1(C)})$ is the quotient map and $\gamma : K_1(C)/\mathrm{Tor}(K_1(C)) \to K_1(B)$ is injective. Moreover, we may assume that $B$ has a unique tracial state, if $A$ has a unique tracial state.*

*Proof.* Without loss of generality, we may assume that $K_1(C)$ is torsion free, by replacing $C$ by $C \otimes \mathcal{U}$.

Let $\tau \in T(A)$ be a tracial state of $A$. There is a unital simple AF-algebra $A_0$ with a unique tracial state and $K_0(A_0) = \rho_A(K_0(A_0)) = \mathbb{D}$, where $\mathbb{D}$ is a subring of $\mathbb{R}$ which is also a countable divisible dense subgroup of $\mathbb{R}$ containing $\tau(K_0(A))$ with $[1_{A_0}] = 1$. There is a unital monomorphism $\psi_1 : A \to A_0$. Therefore, by replacing $j$ by $\psi_1 \circ j$, we may assume that $A = A_0$. We may also assume that $A \otimes \mathcal{U} = A$.

Let $C_1 = C \otimes A$ and $C_2 = C_1 \otimes A \otimes M_2$,...,$C_{n+1} = C_n \otimes A \otimes M_n$, $n = 1, 2, ...,$. We will identify $A$ with $A \otimes M_n$ whenever it is convenient. In particular, $C_{n+1} \cong C_n \otimes A$. Put $A_n = \overbrace{A \otimes A \otimes \cdots \otimes A}^{n}$. Denote $j_1 : C_1 \to A \otimes A$ by $j_1(c \otimes a) = j(c) \otimes a$ for all $a \in A$ and $c \in C$. Define $j_{n+1} : C_{n+1} \to A_{n+2}$ by $j_{n+1}(c \otimes a) = j_n(c) \otimes a$ for all $c \in C_n$ and $a \in A$, $n = 1, 2, ....$

Note that, since $A$ is a unital simple AF-algebra, $K_1(C_1) = K_1(C \otimes \mathcal{U}) \otimes K_0(A)$. Since $K_1(C \otimes \mathcal{U})$ and $K_0(A)$ are torsion free, the map $\varphi_0 : C \otimes \mathcal{U} \to C \otimes \mathcal{U} \otimes A$ defined by $a \to a \otimes 1$ induces an injective homomorphism $(\varphi_0)_{*1}$.

Define $\varphi_1 : C_1 \to C_2$ by

$$\varphi_1(a) = \mathrm{diag}(a, 1 \otimes j_1(a)) \text{ for all } a \in C_1 \text{ and} \tag{e 4.107}$$

$$\varphi_n(a) = \mathrm{diag}(a, \overbrace{1 \otimes j_n(a), 1 \otimes j_n(a), ..., 1 \otimes j_n(a)}^{n-1}) \text{ for all } a \in C_n \tag{e 4.108}$$

$n = 1, 2, ....$ Note that, for each $n > 1$, $C_n \cong C_1 \otimes A_n$. $(\varphi_n)_{*1}$ is injective, $n = 1, 2, ....$ Put $B = \lim_{n\to\infty}(C_n, \varphi_n)$. Then $\varphi_{1,\infty} : C_1 \to B$ is a unital monomorphism. In particular, $(\varphi_{1,\infty})_{*1}|_{K_1(C_1)}$ is injective. Therefore $(\varphi_{1,\infty} \circ \varphi_0)_{*1}$ is injective on $K_1(C \otimes \mathcal{U})$. Since $A$ is a unital simple $C^*$-algebra, it is also easy and standard to check that $B$ is a unital simple $C^*$-algebra.

To verify that $B$ has tracial rank zero, fix any $\epsilon > 0$ and a finite subset $\mathcal{F} \subset B$ and a positive element $a \in B_+ \setminus \{0\}$. Without loss of generality, we may assume that $\mathcal{F} \subset \varphi_{n,\infty}(C_n)$ for some large $n$. There exists a finite subset $\mathcal{G} \subset C_n$ such that $\varphi_{n,\infty}(\mathcal{G}) \supset \mathcal{F}$. Fix an integer $k \geq 1$. Then one may write

$$\varphi_{n,n+k}(x) = \imath(x) \oplus \Phi(x) \text{ for all } x \in C_n, \tag{e 4.109}$$

where $\imath : C_n \to (1-p)C_{n+k}(1-p)$ is a unital monomorphism such that $1 - p = e_{11}$, where $\{e_{ij}\}$ is a system of matrix unit for $M_{n(n+1)\cdots n(+k)}$ and $\Phi : C_n \to F$, where $F$ is a unital AF-subalgebra of $pC_np$ (with $1_F = p$).

Let $F_\infty = \varphi_{n+1,\infty}(F)$. Then $F_\infty$ is an unital AF-algebra with $1_{F_\infty} = \varphi_{n+1,\infty}(p)$. We also have

$$[y, \varphi_{n,\infty}(p)] = 0 \text{ for all } y \in \mathcal{F}. \tag{e 4.110}$$

This, in particular, implies that $B$ satisfies the Popa condition (1) and (2) as in 3.6.2 of [35] (see also [54]). By 3.6.6 of [35], $B$ has the property (SP). Now let $e \in \overline{aBa}$ be a non-zero projection. We will show that we can choose $p$ (by choosing larger $k$ above) so that

$$[1 - p] \leq [e]. \tag{e 4.111}$$

To do this, by replacing $e$ by an equivalent projection, we may assume that $e \in \varphi_{m,\infty}(C_m)$ for some $m \geq 1$. Note that each $\varphi_n$ is injective. So there is a projection $q_0 \in C_m$ such that $\varphi_{m,\infty}(q_0) = e$. Note that

$$\varphi_{m,m+1}(q_0) = \mathrm{diag}(q_0, \overbrace{1 \otimes j_m(q_0), 1 \otimes j_m(q_0), ..., 1 \otimes j_m(q_0)}^{m}) \tag{e 4.112}$$

In particular,

$$[\varphi_{m,\infty}(q_0)] \geq [\varphi_{m+1,\infty}(0, 1 \otimes j_m(q_0), 0, ..., 0)] \tag{e 4.113}$$

In other words, there is a projection $q_1 \in 1 \otimes A \subset C_{m+1}$ such that $[\varphi_{m+1,\infty}(q_1)] \leq [e]$. It then easy to check that, for some sufficiently large $k$, for the above chosen $p$,

$$[1 - p] \leq [\varphi_{m+1,\infty}(q_1)] \leq [e]. \tag{e 4.114}$$

It follows that $B$ has tracial rank zero. Since each $C_n$ satisfies the UCT, so does $B$ (see [59]).

It is also standard to show that $B$ has a unique tracial state if $A$ does (see, for example, 3.7.10 of [35]).

□

**Corollary 4.2.** *Let $C$ be a unital separable $C^*$-algebra in $\mathcal{N}$, let $A \cong A \otimes \mathcal{U}$ be a unital simple AF-algebra with a unique tracial state $\tau$ and with $K_0(A) = \rho_A(K_0(A))$ which is a subring of $\mathbb{R}$ with $[1_A] = 1$ and let $j : C \to A$ be a unital embedding.*

*Then there exists a unital separable simple $C^*$-algebra $B \in \mathcal{N}$ with tracial rank zero, with a unique tracial state, $\rho_B(K_0(B)) = \rho_A(K_0(A))$ and a unital monomorphism $\varphi : C \to B$ satisfying the following:*

$$(\varphi)_{*1} = \gamma \circ \kappa \tag{e 4.115}$$

*where $\kappa : K_1(C) \to K_1(C)/\mathrm{Tor}(K_1(C))$ is the quotient map and $\gamma : K_1(C)/\mathrm{Tor}(K_1(C)) \to K_1(B)$ is injective. Moreover, there is a unital monomorphism $h : B \to A$ and a unital homomorphism with $h_{*0} = \rho_B$ such that*

$$(h \circ \varphi)_{*0} = j_{*0}. \tag{e 4.116}$$

*Proof.* By considering the composition $C \to C \otimes \mathcal{U} \to A \otimes \mathcal{U} \cong A$, we may assume that $K_1(C)$ is torsion free. Since $C$ can be embedded into a unital simple AF-algebra, it has a faithful tracial state. We will keep the notation used in the proof of 4.1. Note that $C_{n+1} \cong C_n \otimes A$ for each $n$ and for each $t \in T(C \otimes A)$, it has the form $t_1 \otimes \tau$, where $t_1$ is a tracial state of $C$ and $\tau \in T(A)$ is the unique tracial state. We will use $\tau$ for the unique tracial state on $A_n$.

The homomorphism $\varphi_n : C_n \to C_{n+1}$ now has the from

$$\varphi_n(a) = \mathrm{diag}(a, \overbrace{1 \otimes j_n(a), 1 \otimes j_n(a), ..., 1 \otimes j_n(a)}^{n-1}) \tag{e 4.117}$$

for all $a \in C_n$. It follows that

$$(t_1 \otimes \tau)(\varphi_n(a)) = \frac{1}{n} t_1(a) + \frac{n-1}{n} \tau(j_n(a)) \text{ for all } a \in C_n. \tag{e 4.118}$$

Thus, we compute that

$$\lim_{n\to\infty} (t_1 \otimes \tau)(\varphi_{k,n}(a)) = \tau(j_k(a)) \text{ for all } a \in C_k. \tag{e 4.119}$$

Since $B$ has a unique tracial state, we compute that

$$\tau_1(\varphi(a)) = \tau(j(a)) \text{ for all } c \in C, \tag{e 4.120}$$

where $\tau_1$ is the unique tracial state of $B$. Since $B \in \mathcal{N}$, by the classification theorem of [40], there is a monomorphism from $B$ to $B_1$ which is a unital simple AH-algebra with a unique tracial state, with $K_0(B_1) = \rho_{B_1}(K_0(B_1)) = \rho_B(K_0(B))$. Then (e 4.116) follows.

□

**4.3.** Let $C_0$ be a unital amenable $C^*$-algebra, let $\Lambda : \mathbb{Z}^k \to Aut(C_0)$ ($k \geq 1$) be a homomorphism, and let $C = C_0 \rtimes_\Lambda \mathbb{Z}^k$. Suppose that $A = A \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state $\tau$ and $K_0(A) = \rho_A(K_0(A))$ which is a subring of $\mathbb{R}$ with $[1_A] = 1$ and that $j : C \to A$ is a unital monomorphism. By 4.2, there exists a unital separable simple $C^*$-algebra $B_0 \in \mathcal{N}$ with tracial rank zero, with a unique tracial state and with $\rho_B(K_0(B_0)) = \rho_A(K_0(A))$, and there exists a unital monomorphism $j_1 : C \to B_0$ satisfying the following:

$$(j_1)_{*1} = \gamma \circ \kappa \tag{e 4.121}$$

where $\kappa : K_1(C) \to K_1(C)/\mathrm{Tor}(K_1(C))$ is the quotient map and $\gamma : K_1(C)/\mathrm{Tor}(K_1(C)) \to K_1(B_0)$ is injective.

Note also in the proof of 4.1 $K_1(C_n) = K_1(C) \otimes K_0(A)$ and the homomorphism $\varphi_n : K_1(C_n) \to K_1(C_{n+1})$ is the identity map. So $K_1(B_0) = K_1(C) \otimes K_0(A)$.

Moreover, there is a unital monomorphism $h : B_0 \to A$ with $h_{*0} = \rho_{B_0}$ such that

$$(h \circ j_1)_{*0} = j_{*0}. \tag{e 4.122}$$

Note that $K_0(A)$ is also divisible. It follows that

$$[h \circ j_1] = [j] \text{ in } KK(C, A) \text{ and } \tau(h \circ j_1) = \tau \circ j. \tag{e 4.123}$$

There exists (see 2.8 and [30]) a homomorphism $R_{j,h\circ\varphi} : K_1(C) \to Aff(T(A)) = \mathbb{R}$. Then $R_{j,h\circ j_1}(K_1(C))$ is a countable subgroup of $\mathbb{R}$. Fix a countable subring $\mathbb{D} \subset \mathbb{R}$ which contains $\rho_A(K_0(A))$ and $R_{j,h\circ j_1}(K_1(C))$ which also has the property that $\mathbb{D}\mathbb{Q} = \mathbb{D}$. Let $B = B(j, A)$ be a unital simple AF-algebra with unique tracial state and $K_0(B) = \rho_B(K_0(B)) = \mathbb{D}$ and with $[1_B] = 1$. There is a unital embedding $j_2 : A \to B$ such that $(j_2)_{*0} = \mathrm{id}_{\rho_A(K_0(A))}$. We may assume that $A \subset B$.

An important fact is that, if we regard $j : C \to B$ and $h \circ j_1 : C \to B$, then

$$R_{j,h\circ j_1}(K_1(C)) \subset \mathbb{D} = \rho_B(K_0(B)). \tag{e 4.124}$$

**Theorem 4.4.** *Let $C_0$ be a unital AH-algebra, $C = C_0 \rtimes_\alpha \mathbb{Z}$ for some $\alpha \in Aut(C_0)$, let $A \cong A \otimes \mathcal{U}$ be a unital simple AF-algebra with a unique tracial state and with $K_0(A) = \rho_A(K_0(A))$ which is a subring of $\mathbb{R}$ and let $j : C \to A$ be a unital monomorphism. Let $B = B(j, A)$ and $\mathbb{D} = K_0(B)$ be as in* 4.3.

*Let $\epsilon > 0$, let $\mathcal{F} \subset C$ be a finite subset and let $\mathcal{P} \subset K_1(C)$. There is $\delta > 0$ and a finite subset $\{y_1, y_2, ..., y_m\} \subset K_1(C)$ satisfying the following:*

*If $\beta : K_1(C) \to K_0(A)$ is a homomorphism with*

$$\rho_A((\beta(y_j))(\tau) < \delta, \ \ j = 1, 2, ..., m \tag{e 4.125}$$

*where $\tau$ is the unique tracial state of $A$, then there exists a unitary $u \in B$ such that*

$$\|[\psi(a),\, u]\| < \epsilon \ \textit{for all} \ a \in \mathcal{F} \ \text{and} \ \mathrm{bott}_1(\psi,\, u)|_{\mathcal{P}} = \beta'|_{\mathcal{P}} \tag{e 4.126}$$

*where $\psi : C \to B \otimes \mathcal{U} \to B$ is defined by $\psi(c) = j(c) \otimes 1_{\mathcal{U}}$ for $c \in C_0$ and $\psi(u_\alpha) = j(u_\alpha) \otimes \imath(u_\sigma)$ and $\beta' = (j_2)_{*0} \circ \beta$ ($j_2$ is as in 4.3).*

*Proof.* Since $K_0(A)$ is torsion free, $\beta(\mathrm{Tor}(K_1(C))) = \{0\}$. Therefore, there exists $\beta_1 : K_1(C)/\mathrm{Tor}(K_1(C)) \to K_0(A)$ such that

$$\beta = \beta_1 \circ \kappa, \tag{e 4.127}$$

where $\kappa : K_1(C) \to K_1(C)/\mathrm{Tor}(K_1(C))$ is the quotient map.

Let $B_0$ be as in 4.3. By the classification of amenable simple $C^*$-algebras of tracial rank zero ([40]), $B_0$ is a unital simple AH-algebra of real rank zero. Write $B_0 = \lim_{n\to\infty}(C_n, \varphi_n)$, where each $C_n$ has the form $P_n M_{k(n)}(C(X_n))P_n$ and where $X_n$ is a finite CW complex and $P_n \in M_{k(N)}(C(X_n))$ is a projection. Moreover, let $\gamma$ and $j$ be as in 4.3.

Identify $K_0(A)$ with $\rho_A(K_0(A)) \subset \mathbb{R}$. We define $\gamma_1 : K_1(B_0) = K_1(C)\otimes\rho_A(K_0(A)) \to K_0(A)$ by

$$\gamma_1(g\otimes r) = r\beta(g) \text{ for all } r \in \rho_A(K_0(A)) \text{ and } g \in K_1(C). \tag{e 4.128}$$

Then

$$\gamma_1 \circ \gamma = \beta_1. \tag{e 4.129}$$

Furthermore, for any $x \in K_1(B_0)$, there are $g_1, g_2, ..., g_l \in K_1(C)/\mathrm{Tor}(K_1(C))$ and real numbers $r_1, r_2, ..., r_l \in \rho_A(K_0(A))$ such that

$$\gamma_1(x) = \sum_{i=1}^{l} r_i\beta_1(g_i). \tag{e 4.130}$$

Since $B_0$ satisfies the Universal Coefficient Theorem, there is a homomorphism $\tilde{\beta} \in Hom_\Lambda(\underline{K}(B_0), \underline{K}(SB))$ such that

$$\tilde{\beta}|_{K_1(B_0)} = \gamma_1. \tag{e 4.131}$$

Without loss of generality, we may assume that $j_1(\mathcal{F}) \subset \varphi_{n,\infty}(C_n)$ and $j_1(\mathcal{P}) \subset [\varphi_{n,\infty}](\underline{K}(C_n))$.

Let $\delta_0 > 0$ (in place of $\delta$) and $k(n) \geq n$ and $x_1, x_2, ..., x_k$ be a set of generators for $K_1(C_{k(n)})$ required by Lemma 7.5 of [49] corresponding to $\epsilon/2$ and $\mathcal{F}_1 = h(\mathcal{F})$. By (e 4.130), there are $y_1, y_2, ..., y_m \in K_1(C)$ such that

$$\gamma_1(x_j) = \sum_{i=1}^{m} r_{i,j}\beta(y_i), \;\; j = 1, 2, ..., k. \tag{e 4.132}$$

Therefore there exists $\delta > 0$ such that

$$\rho_B(\beta(y_j))(\tau) < \delta, \;\; j = 1, 2, ..., m, \tag{e 4.133}$$

implies

$$\rho_B(\gamma_1(x_i))(\tau) < \delta_0, \;\; i = 1, 2, ..., k. \tag{e 4.134}$$

We now choose the above $\delta$ and $y_1, y_2, ..., y_m$ and assume (e 4.125) holds.

Thus, by 7.5 of [49], there is a unitary $u_0 \in A$ such that

$$\|[h,\, u_0]\| < \epsilon/2 \text{ for all } a \in \mathcal{F}_1 \text{ and } \mathrm{bott}_1(h, u_0)|_{\mathcal{P}} = \bar{\beta}|_{\mathcal{P}}. \tag{e 4.135}$$

Viewing $A$ as a unital $C^*$-subalgebra of $B$, by 4.3,

$$[h\circ j_1] = [j] \text{ in } KK(C, B), \;\; \tau\circ h\circ j_1 = \tau\circ j \text{ and} \tag{e 4.136}$$

$$R_{j,h\circ j_1}(K_1(C)) \subset \rho_B(K_0(B)). \tag{e 4.137}$$

Thus, by 3.6, there is a unitary $w \in U(B)$ such that

$$\mathrm{ad}\, w\circ\psi_1 \approx_{\epsilon/2} \psi \text{ on } \mathcal{F}, \tag{e 4.138}$$

where $\psi_1(c) = h\circ j_1(c)\otimes 1$ and $\psi_1(u_\alpha) = h\circ j_1(u_\alpha)\otimes\imath(u_\sigma)$. Choose $u = w^*u_0w$. Then

$$\|[\psi(a),\, u]\| < \epsilon \text{ for all } a \in \mathcal{F} \text{ and } \mathrm{bott}_1(\psi, u)|_{\mathcal{P}} = \beta|_{\mathcal{P}}. \tag{e 4.139}$$

□

# 5 Homotopy lemmas

The next follows immediately from Theorem 3.9 of [43]. Several versions of the following have been appeared.

**Theorem 5.1.** *Let $C$ be a unital separable amenable $C^*$-algebra. For any $\epsilon > 0$ and any finite subset $\mathcal{F} \subset C$, there exists $\delta > 0$, a finite subset $\mathcal{G} \subset C$, a finite subset $\mathcal{Q} \subset \underline{K}(C)$ and an integer $n \geq 1$ satisfying the following:*

*For any unital simple $C^*$-algebra $A$ of tracial rank zero and any two unital $\delta$-$\mathcal{G}$-multiplicative contractive completely positive linear maps $L_1, L_2 : C \to A$ and any unital monomorphism $h : C \to A$, if*

$$[L_1]|_{\mathcal{Q}} = [L_2]|_{\mathcal{Q}}, \tag{e 5.140}$$

*then there exists a unitary $U \in M_{n+1}(A)$ such that*

$$\mathrm{ad}\, U \circ \bar{L}_1 \approx_\epsilon \bar{L}_2 \ \ on \ \mathcal{F}, \tag{e 5.141}$$

*where*

$$\bar{L}_i(a) = \mathrm{diag}(L_1(a), \overbrace{h(a), h(a), ..., h(a)}^{n})$$

*for all $a \in C$ and $i = 1, 2$.*

*Proof.* We will apply Theorem 1.2 of [24]. By Theorem 3.9 of [43] and the assumption that there exists a unital embedding $h : C \to A$, $C$ satisfies property (P) of Remark 1.1 of [24]. Thus Theorem 1.2 of [24] holds for $C$ (in place $A$) and $A$ (in place of $B$). Since in this case $A$ has tracial rank zero, weakly unperforated $K_0(A)$, stable rank one and real rank zero, the theorem follows ( see also the proof of Theorem 3.1 of [24]).

□

**Lemma 5.2.** *Let $C_0$ be a unital AH-algebra, $C = C_0 \rtimes_\alpha \mathbb{Z}$ for some $\alpha \in Aut(C_0)$ and let $A \cong A \otimes \mathcal{U}$ be a unital simple AF-algebra with a unique tracial state $\tau$ and with $K_0(A) = \rho_A(K_0(A))$. Let $j : C \to A$ be a unital monomorphism, $\epsilon > 0$, $\mathcal{F} \subset C_0$ be a finite subset and $\mathcal{P} \subset K_1(C_0)$ be a finite subset. There is $\delta > 0$, a finite subset $\mathcal{G} \subset C_0$, and integer $K$ and a finite subset $\mathcal{Q} \subset \underline{K}(C)$ satisfying the following:*

*Suppose that $v \in U(A)$ is a unitary such that*

$$\|[j(a),\, v]\| < \delta \ \ for \ \ all \ \ a \in \mathcal{G} \ \text{ and} \tag{e 5.142}$$

$$[L]|_{\mathcal{Q}} = [j]|_{\mathcal{Q}}, \tag{e 5.143}$$

*where $L : C \to A$ is a contractive completely positive linear map such that*

$$\|L(\sum_{k=-K}^{K} f_k u_\alpha^k) - \sum_{k=-K}^{K} j(f_k)(j(u_\alpha)v)^k\| < \delta \tag{e 5.144}$$

*for all $f_k \in \mathcal{G}$. Then there exists a unitary $w \in U(A)$ such that*

$$\|[j(a),\, w]\| < \epsilon \ \ for \ \ all \ \ a \in \mathcal{F} \ \text{ and } \ \mathrm{bott}_1(j|_{C_0}, j(u_\alpha)^* w^* j(u_\alpha) v w)|_{\mathcal{P}} = 0. \tag{e 5.145}$$

*Proof.* Let $\epsilon > \epsilon_0 > 0$ and $\mathcal{F} \subset \mathcal{F}_1 \subset C_0$ be a finite subset. We assume that

$$\mathrm{bott}_1(h,\, v')|_{\mathcal{P}} \tag{e 5.146}$$

is well defined for any unital homomorphism $h$ from $C_0$ and any unitary $v'$ provided that $\|[h(a),\, v']\| < \epsilon_0$ for all $a \in \mathcal{F}_1$.

Let $\mathcal{P} = \{x_1, x_2, ..., x_k\}$.

We write $C_0 = \lim_{n\to\infty}(C_n, \varphi_n)$, where each $C_n$ has the form $P_nM_{r(n)}(C(X_n))P_n$, where $X_n$ is a finite CW complex and $P_n \in M_{r(n)}(C(X_n))$ is a projection. We may assume that $\mathcal{P} \subset [\varphi_{n,\infty}](K_1(C_n))$. Let $\eta > 0$ (in place of $\delta$), $k(n) \ge n$ and $y_1, y_2, ..., y_m \in K_1(C_{k(n)})$ be a set of generators as required by Lemma 7.3 of [49] for $\{x_1, x_2, ..., x_k\}$, $\epsilon_0/2$ and $\mathcal{F}_1$. Denote $z_j = (\varphi_{k(n),\infty})_{*1}(y_j)$, $j = 1, 2, ..., m$. We may assume that $\mathcal{P} \subset \{z_1, z_2, ..., z_m\}$.

Let $\mathcal{P}_1' \subset \underline{K}(C_0)$ be a finite subset containing $z_1, z_2, ..., z_m$ and $\mathcal{P}_1 = \mathcal{P}_1' \cup [\alpha](\mathcal{P}_1')$.

Let $\epsilon_1 > 0$ and $\mathcal{F}_2 \subset C_0$ be a finite subset such that

$$\mathrm{bott}_1(h',\, v')|_{\mathcal{P}_1} \tag{e 5.147}$$

is well defined for any unital homomorphism $h'$ from $C_0$ and any unitary $v'$ provided that $\|[h'(a),\, v']\| < \epsilon_1$ for all $a \in \mathcal{F}_2$. We may also

$$\mathrm{bott}_1(h'', v')|_{\mathcal{P}_1} = \mathrm{bott}_1(h', v')|_{\mathcal{P}_1} = \mathrm{Bott}(h', v'')|_{\mathcal{P}_1}, \tag{e 5.148}$$

provided that $h'' \approx_{\epsilon_1} h'$ on $\mathcal{F}_2$ and $\|v' - v''\| < \epsilon_1$. We may further assume that $\epsilon_1 < \epsilon_0/2$ and $\mathcal{F}_1 \subset \mathcal{F}_2$.

To simplify notation, we may also assume that

$$\rho_D(\mathrm{bott}_1(h', v'))(z_j)(t) < \eta, \;\; j = 1, 2, ..., m \tag{e 5.149}$$

for any unital homomorphism $h' : C \to D$, any unitary $v' \in D$ and unital $C^*$-algebra $D$ and tracial state $t \in T(D)$, provided that

$$\|[h'(a),\, v']\| < \epsilon_1 \;\text{ for all }\; a \in \mathcal{F}_2. \tag{e 5.150}$$

Put $\mathcal{G}_1 = \mathcal{F}_2 \cup \{u_\alpha\}$.

Let $\delta_1 > 0$ ( in place of $\delta$) a finite subset $\mathcal{G}_2 \subset C$ (in place of $\mathcal{G}$) be a finite subset, $\mathcal{Q} \subset \underline{K}(C)$ be a finite subset and $N \ge 1$ be an integer required by 5.1 for $\epsilon_1/2$ and $\mathcal{G}_1$.

There is $\delta$, a finite subset $\mathcal{G} \subset C$ and an integer $K \ge 1$ such that $L$ is $\delta_1$-$\mathcal{G}_1$-multiplicative if (e 5.142) and (e 5.144) hold.

Suppose (e 5.142), (e 5.143) and (e 5.144) hold.

It follows from 5.1 that there exists a unitary $U \in M_{N+1}(A)$ such that

$$\mathrm{ad}\, U \circ \bar{L} \approx_{\epsilon_1/2} \bar{j} \;\text{ on }\; \mathcal{G}_1, \tag{e 5.151}$$

where

$$\bar{L}(a) = \mathrm{diag}(L(a), \overbrace{j(a), j(a), ..., j(a)}^{N}) \;\text{ and} \tag{e 5.152}$$

$$\bar{j}(a) = \mathrm{diag}(\overbrace{j(a), j(a), ..., j(a)}^{N+1}) \tag{e 5.153}$$

for all $a \in C$. It follows that

$$\|[\bar{j}(a),\, U]\| < \epsilon_1/2 \;\text{ for all }\; a \in \mathcal{F}_2. \tag{e 5.154}$$

Put $V = \mathrm{diag}(\overbrace{v, v, ..., v}^{N+1})$. By (e 5.151),

$$\|1 - \bar{j}(u_\alpha)^* U^* \bar{j}(u_\alpha) V U\| < \epsilon_1/2. \tag{e 5.155}$$

It follows that

$$\mathrm{bott}(\bar{j}|_{C_0},\, \bar{j}(u_\alpha)^* U^* \bar{j}(u_\alpha) V U)|_{\mathcal{P}_1} = 0. \tag{e 5.156}$$

Thus (by (e 5.148)),

$$\begin{aligned} 0 = \;& \mathrm{bott}_1(\bar{j}|_{C_0},\, V)|_{\mathcal{P}_1} + \mathrm{bott}_1(\bar{j}|_{C_0},\, V^* \bar{j}(u_\alpha)^* U^* \bar{j}(u_\alpha) V U) && \text{(e 5.157)} \\ = \;& \mathrm{bott}_1(\bar{j}|_{C_0},\, V)|_{\mathcal{P}_1} + \mathrm{bott}_1(\bar{j}|_{C_0},\, \bar{j}(u_\alpha)^* U^* \bar{j}(u_\alpha) V U V^*)|_{\mathcal{P}_1} && \text{(e 5.158)} \\ = \;& \mathrm{bott}_1(\bar{j}|_{C_0}, V)|_{\mathcal{P}_1} + \mathrm{bott}_1(\bar{j}|_{C_0},\, \bar{j}(u_\alpha)^* U^* \bar{j}(u_\alpha))|_{\mathcal{P}_1} + \mathrm{bott}_1(\bar{j}|_{C_0}, V U V^*)|_{\mathcal{P}_1} && \text{(e 5.159)} \\ = \;& \mathrm{bott}_1(\bar{j}|_{C_0}, V)|_{\mathcal{P}_1} + \mathrm{bott}_1(\bar{j} \circ \alpha^{-1}|_{C_0},\, U^*)|_{\mathcal{P}_1} + \mathrm{bott}_1(\bar{j}|_{C_0}, U)|_{\mathcal{P}_1} && \text{(e 5.160)} \end{aligned}$$

Let $G_1$ be a subgroup of $K_1(C)$ generated by $\mathcal{P}_1$. Since $K_0(A)$ is divisible, there are homomorphisms $\lambda_1, \gamma : K_1(C_0) \to K_0(A)$ which extend $\mathrm{bott}_1(\bar{j}|_{C_0},\, V)$ and $\mathrm{bott}_1(\bar{j}|_{C_0},\, U)$, respectively.

It follows from (e 5.160) that

$$\lambda_1|_{G_1} = -\gamma|_{G_1} + \gamma \circ \alpha_{*1}^{-1}|_{G_1}. \tag{e 5.161}$$

It should be note that

$$(\lambda_1)|_{G_1} = (N+1)\mathrm{bott}_1(j|_{C_0}, v)|_{G_1}. \tag{e 5.162}$$

Define $\gamma_1 : K_1(C_0) \to K_0(A)$ by $\gamma_1(x) = -\frac{1}{N+1}\gamma(x)$ for all $x \in K_1(C_0)$.

By (e 5.149) and applying Lemma 7.3 of [49], there is a unitary $w \in A$ such that

$$\|[j(a),\, w]\| < \epsilon_0/2 \;\text{ and }\; \mathrm{bott}_1(j,\, w)(x_j) = \gamma_1(x_j), \;\; j = 1, 2, ..., k. \tag{e 5.163}$$

We then compute (using (e 5.161) among other things) that

$$\begin{aligned} & \mathrm{bott}_1(j,\, j(u_\alpha^*) w^* j(u_\alpha) v w)(x_j) && \text{(e 5.164)} \\ & \quad = \; \mathrm{bott}_1(j, v)(x_j) + \mathrm{bott}_1(j,\, v^* j(u_\alpha)^* w^* j(u_\alpha) v w)(x_j) && \text{(e 5.165)} \\ & \quad = \; \mathrm{bott}_1(j,\, v)(x_j) + \mathrm{bott}_1(j,\, j(u_\alpha)^* w^* j(u_\alpha) v w v^*)(x_j) && \text{(e 5.166)} \\ & \quad = \; \mathrm{bott}_1(j, v)(x_j) + \mathrm{bott}_1(j,\, j(u_\alpha)^* w^* j(u_\alpha))(x_j) + \mathrm{bott}_1(j, w v w^*)(x_j) && \text{(e 5.167)} \\ & \quad = \; \mathrm{bott}_1(j, v)(x_j) + \mathrm{bott}_1(j \circ \alpha^{-1},\, w^*)(x_j) + \mathrm{bott}_1(j, w)(x_j) && \text{(e 5.168)} \\ & \quad = \; \frac{1}{N+1}(\lambda_1)(x_j) + (-\gamma_1 \circ (\alpha_{*1}^{-1}(x_j) + \gamma_1(x_j)) && \text{(e 5.169)} \\ & \quad = \; (\frac{1}{N+1})(\lambda_1 - \gamma + \gamma \circ \alpha_{*1}^{-1})(x_j) = 0, && \text{(e 5.170)} \end{aligned}$$

$j = 1, 2, ..., k$.

It follows that

$$\mathrm{bott}_1(j_{C_0},\, j(u_\alpha^*) w^* j(u_\alpha) v w)|_{\mathcal{P}} = 0. \tag{e 5.171}$$

□

**Theorem 5.3.** *Let $C$ be a unital AH-algebra and let $\alpha \in Aut(C)$ be an automorphism and let $A \cong A \otimes \mathcal{U}$ be a unital simple AF-algebra with a unique tracial state $\tau$ and $\rho_A(K_0(A)) = K_0(A)$. Let $h : (C \rtimes_\alpha \mathbb{Z}) \otimes C(\mathbb{T}) \to A$ be a unital monomorphism.*

*For any $\epsilon > 0$ and a finite subset $\mathcal{F} \subset C \rtimes_\alpha \mathbb{Z}$, there exists $\delta > 0$, $\eta > 0$, a finite subset $\mathcal{G} \subset C \rtimes_\alpha \mathbb{Z}$, a finite subset $\bar{\mathcal{G}} \subset C \otimes C(\mathbb{T})$ and a finite subset $\mathcal{P} \subset K_1(C \rtimes_\alpha \mathbb{Z})$ satisfying the following:*

*Suppose that there is a unitary $v \in U(A)$ and contractive completely positive linear map $L : C \otimes C(\mathbb{T}) \to A$ such that*

$$\|[h(a),\, v]\| < \delta \ \textit{for all} \ a \in \mathcal{G}, \mathrm{bott}_1(h, v)|_{\mathcal{P}} = 0 \ \text{and} \tag{e 5.172}$$

$$\tau \circ h|_{C\otimes C(\mathbb{T})} \approx_\eta \tau \circ L \ \textit{on} \ \bar{\mathcal{G}}, \tag{e 5.173}$$

*and*

$$L \approx_\delta h \ \textit{on} \ \mathcal{G} \ \text{and} \ L(1 \otimes z) \approx_\delta v \tag{e 5.174}$$

*and where $z \in C(\mathbb{T})$ is the identity function on the unit circle. Then there exists a unitary $W \in A \otimes \mathcal{U}$ such that*

$$\|[h^{(1)}(c),\, W]\| < \epsilon \ \text{for all} \ c \in \mathcal{F} \ \text{and} \ W^*(v \otimes 1_{\mathcal{U}})W \approx_\epsilon h(1 \otimes z) \otimes 1_{\mathcal{U}}, \tag{e 5.175}$$

*where $h^{(1)}(c) = h(c) \otimes 1_{\mathcal{U}}$ for all $c \in C$ and $h^{(1)}(u_\alpha) = h(u_\alpha) \otimes \imath(u_\sigma)$.*

*Proof.* Let $\epsilon > 0$ and $\mathcal{F} \subset C \rtimes_\alpha \mathbb{Z}$. Without loss of generality, we may assume that $\mathcal{F} = \mathcal{F}_0 \cup \{u_\alpha\}$ for some finite subset $\mathcal{F}_0 \subset C$. Define $C_1 = C \otimes C(\mathbb{T})$. Define $\alpha' \in Aut(C_1)$ by $\alpha'(c \otimes 1_{C(\mathbb{T})}) = \alpha(c) \otimes 1_{C(\mathbb{T})}$ and $\alpha'(1_C \otimes f) = 1_C \otimes f$ for all $f \in C(\mathbb{T})$. In other words, $C_1 \rtimes_{\alpha'} \mathbb{Z} \cong (C \rtimes_\alpha \mathbb{Z}) \otimes C(\mathbb{T})$. Define $\mathcal{F}' = \mathcal{F}_0 \cup \{1 \otimes z\}$.

To apply 3.2, let $\delta_1 > 0$ (in place of $\delta$), $\mathcal{G}_1 \subset C_1$ (in place of $\mathcal{G}$) be a finite subset and $\mathcal{P}_1' \subset \underline{K}(C_1)$ (in place of $\mathcal{P}$) be a finite subset required by 3.2 for $\epsilon/2$ and $\mathcal{F}'$. We may assume that $\delta_1 < \epsilon/2$ and $\mathcal{G}_1 \supset \mathcal{F}'$. Since $K_1(A) = \{0\}$ and $K_0(A)$ is torsion free and divisible, we may assume that $\mathcal{P}_1' = \boldsymbol{\beta}(\mathcal{P}_1)$ for some finite subset $\mathcal{P}_1 \subset K_1(C_1)$.

Let $\delta_2 > 0$ (in place of $\delta$), $\mathcal{G}_2 \subset C_1$ be a finite subset, $K \geq 1$ be an integer and $\mathcal{Q} \subset \underline{K}(C_1 \rtimes_{\alpha'} \mathbb{Z})$ be a finite subset required by 5.2 for $\delta_1/2$, $\mathcal{P}_1$ and $\mathcal{G}_1$. Since $K_0(A)$ is torsion free and divisible and $K_1(A) = \{0\}$, we may assume that $\mathcal{Q} \subset K_0(C_1 \rtimes_{\alpha'} \mathbb{Z}) = K_0((C \rtimes_\alpha \mathbb{Z}) \otimes C(\mathbb{T}))$. We may assume that $\mathcal{Q} = \mathcal{P}_2 \cup \boldsymbol{\beta}(\mathcal{P}_3)$, where $\mathcal{P}_2 \subset K_0(C \rtimes_\alpha \mathbb{Z})$ and $\mathcal{P}_3 \subset K_1(C \rtimes_\alpha \mathbb{Z})$ are finite subsets. We may assume that $\mathcal{G}_2 \supset \mathcal{G}_1$.

By 2.14, choosing $0 < \delta_2 < \delta_2'$ and a finite subset $\mathcal{G}_2' \supset \mathcal{G}_2$ so that there exits a unital contractive completely positive linear map $L_1 : C_1 \rtimes_{\alpha'} \mathbb{Z} \to A \otimes \mathcal{U}$ such that

$$\|L_1(\sum_{k=-K}^{K} g_k u_\alpha^k) - \sum_{k=-K}^{K} h(g_k)(h(u_\alpha)v')^k\| < \delta_2/2 \tag{e 5.176}$$

for all $g_k \in \mathcal{G}_2$ and for any $v'$ such that $\|[h(a),\, v']\| < \delta_2'$ for all $a \in \mathcal{G}_2'$. We may assume that $\mathcal{G}_2' = \mathcal{G}_2'' \otimes \{1_{C(\mathbb{T})}, z\}$ for some finite subset $\mathcal{G}_2'' \subset C$. Without loss of generality, we may also assume that

$$[L_1]|_{\mathcal{Q}} = [L_1']|_{\mathcal{Q}} \tag{e 5.177}$$

provided that $L_1, L_1'$ both satisfy(e 5.176) $L_1 \approx_{\delta_2} L_1'$ on $\mathcal{G}_2 \cup \{u_\alpha\}$.

Define $\tau_0 = \tau \circ h$ on $C_1$.

Let $0 < \delta_3 < \delta_2$, $\eta > 0$ (in place of $\sigma$), $\bar{\mathcal{G}} \subset C_1$ (in place of $\mathcal{G}$) be a finite subset and $\mathcal{P}_4 \subset \underline{K}(C_1)$ be a finite subset required by 2.13 for $\min\{\delta_2/4, \epsilon/4)\}$ and $\mathcal{G}_2'$ (with $\tau_0$ above). Since

$K_0(A)$ is torsion free and divisible and $K_1(A)=0$, we may assume that $\mathcal{P}_4\subset K_0(C_1)$. Without loss of generality, (by choosing smaller $\delta_3$, for example), we may assume that $\bar{\mathcal{G}}=\mathcal{G}_3'\otimes\mathcal{G}_3''$, where $\{1_C\}\subset\mathcal{G}_3'\subset C$ and $\{1_{C(\mathbb{T})}, z\}\subset\mathcal{G}_3''\subset C(\mathbb{T})$ are finite subsets, and we may assume that $\mathcal{P}_4=\mathcal{P}_5\cup\boldsymbol{\beta}(\mathcal{P}_6)$, where $\mathcal{P}_5\subset K_0(C)$ and $\mathcal{P}_6\subset K_1(C)$ are finite subsets.

Without loss of generality, we may assume that

$$[L]|_{\mathcal{P}_5}=[L']|_{\mathcal{P}_5} \tag{e 5.178}$$

provided that $L, L'$ are $\delta_3$-$\mathcal{G}_3'$-multiplicative contractive completely positive linear maps from $C_1$ and $L\approx_{\delta_3}L'$ on $\mathcal{G}_3'$.

Denote by $j_0: C\to C\rtimes_\alpha\mathbb{Z}$ the natural embedding. Put $\mathcal{P}=(j_0)_{*1}(\mathcal{P}_6)\cup\mathcal{P}_3$. Let $\eta_1>0$ such that $\eta_1<\min\{\delta_3/4,\delta_2/4,\epsilon/2,\delta_1/4,\eta\}$.

It follows from 2.14 that there is $\delta_4>0$, a finite subset $\mathcal{G}_4\subset C$ and a unital $\eta_1$-$\bar{\mathcal{G}}$-multiplicative contractive completely positive linear map $L': C_1\to A$ such that

$$L'(c\otimes f)\approx_{\eta_1} h(c)f(v)\ \text{ for all }\ c\in\mathcal{G}_2'\ \text{ and }\ f\in\mathcal{G}_2'', \tag{e 5.179}$$

provided that $\|[h(a),\, v]\|<\delta_4$ for all $a\in\mathcal{G}_4$. We may assume that $\mathcal{G}_4\supset\mathcal{G}_2''\cup\mathcal{G}_2'''$.

Put $\mathcal{G}=\mathcal{G}_4\cup\{u_\alpha\}$.

Put $\delta=\min\{\delta_4/2,\eta_1/2\}$. Suppose that (e 5.172), (e 5.173) and (e 5.174) hold for the above $\delta$, $\eta$, $\mathcal{G}$, $\bar{\mathcal{G}}$ and $\mathcal{P}$ (and for some $v$ and $L$). Therefore, we may assume that $L$ is an $\eta_1$-$\mathcal{G}_2$-multiplicative contractive completely positive linear map satisfying (e 5.179) (replacing $L'$ by $L$). We still have that

$$\tau\circ L\approx_\eta\tau_0\ \text{ on }\ \bar{\mathcal{G}}. \tag{e 5.180}$$

Note that

$$K_0(C_1)=K_0(C)\oplus K_1(C). \tag{e 5.181}$$

We have (by (e 5.178))

$$[L]|_{\mathcal{P}_5}=h_{*0}|_{\mathcal{P}_5}. \tag{e 5.182}$$

By (e 5.178) and (e 5.172) (and the choice of $\mathcal{P}$), we may also assume that

$$[L]|_{\boldsymbol{\beta}(\mathcal{P}_6)}=0. \tag{e 5.183}$$

Since $\ker\rho_A=\{0\}$,

$$h_{*0}|_{\boldsymbol{\beta}(K_1(C))}=0. \tag{e 5.184}$$

Therefore

$$[h]|_{\mathcal{P}_4}=[L]|_{\mathcal{P}_4}. \tag{e 5.185}$$

Moreover, by (e 5.180),

$$\tau\circ h=\tau_0\approx_\eta\tau\circ L\ \text{ on }\ \bar{\mathcal{G}}. \tag{e 5.186}$$

By applying 2.13, there exists a unitary $w_0\in U(A)$ such that

$$\mathrm{ad}\, w_0\circ L\approx_{\min\{\delta_2/4,\epsilon/2)\}} h\ \text{ on }\ \mathcal{G}_2'. \tag{e 5.187}$$

In particular,

$$w_0^*vw_0 \approx_{\min\{\delta_2/4,\epsilon/2\}} h(1\otimes z) \ \text{ and } \ \|[h(c),\, w_0]\| < \min\{\delta_2/4, \epsilon/2\} \ \text{ for all } \ c\in \mathcal{G}_2''. \tag{e 5.188}$$

Define $V = h(u_\alpha)^*w_0^*h(u_\alpha)w_0$. Then

$$\|[h(a),\, V]\| < \min\{\epsilon/2, \delta_2/4\} \ \text{ for all } \ a\in \mathcal{G}_2'. \tag{e 5.189}$$

By (e 5.189), there exists a unital contractive completely positive linear map $L_1 : C_1\rtimes_{\alpha'}\mathbb{Z}\to A\otimes\mathcal{U}$ which satisfies (e 5.176) with $v' = V$. Therefore

$$L_1 \approx_{\delta_2} h \ \text{ on } \ \mathcal{G}_2'' \ \text{ and } \ L_1(u_\alpha) \approx_{\delta_2} h(u_\alpha)V = w_0^*h(u_\alpha)w_0. \tag{e 5.190}$$

Note that

$$[L_1]|_{\mathcal{P}_2} = [h]|_{\mathcal{P}_2}. \tag{e 5.191}$$

By the assumption (e 5.172) and by (e 5.188), we have

$$\begin{aligned} \mathrm{bott}_1(\mathrm{ad}\, w_0\circ h,\, h(1\otimes z))|_{\mathcal{P}_3} &= \mathrm{bott}_1(\mathrm{ad}\, w_0\circ h,\, w_0^*vw_0)|_{\mathcal{P}_3} && \text{(e 5.192)}\\ &= \mathrm{bott}_1(h,\, v)|_{\mathcal{P}_3} = 0. && \text{(e 5.193)}\end{aligned}$$

It follows from (e 5.188), (e 5.190), (e 5.177) and (e 5.193) that

$$[L_1]|_{\boldsymbol{\beta}(\mathcal{P}_3)} = 0. \tag{e 5.194}$$

Since $\ker\rho_A(A) = \{0\}$, we have

$$[L_1]|_{\boldsymbol{\beta}(\mathcal{P}_3)} = 0 = [h]|_{\boldsymbol{\beta}(\mathcal{P}_3)}. \tag{e 5.195}$$

Thus

$$[L_1]|_{\mathcal{Q}} = [h]|_{\mathcal{Q}}. \tag{e 5.196}$$

Therefore, by 5.2, there is a unitary $w_1\in U(A\otimes\mathcal{U})$ such that

$$\|[h(a),\, w_1]|\| < \delta_1/2 \ \text{ for all } \ a\in\mathcal{G}_1 \ \text{ and } \ \mathrm{Bott}(h,\, h(u_\alpha)^*w_1^*h(u_\alpha)Vw_1)|_{\mathcal{P}_1} = 0. \tag{e 5.197}$$

Note that $h(u_\alpha)V = w_0^*h(u_\alpha)w_0$. It follows from 3.2 that there exists $w_2\in U(A\otimes\mathcal{U})$ such that

$$\begin{aligned} w_2^*(h(a)\otimes 1_{\mathcal{U}})w_2 &\approx_{\epsilon/2} h(a)\otimes 1_{\mathcal{U}} \ \text{ for all } \ a\in\mathcal{F}' \ \text{ and} && \text{(e 5.198)}\\ w_2^*(w_1^*(w_0^*h(u_\alpha)w_0)w_1)\otimes\imath(u_\sigma))w_2 &\approx_{\epsilon/2} h(u_\alpha)\otimes\imath(u_\sigma). && \text{(e 5.199)}\end{aligned}$$

Define $W = (w_0w_1\otimes 1_{\mathcal{U}})w_2$. Then, by (e 5.188), (e 5.198) and (e 5.199),

$$\|[h^{(1)}(c),\, W]\| < \epsilon \ \text{ for all } \ c\in\mathcal{F} \ \text{ and } \ W^*(v\otimes 1_{\mathcal{U}})W \approx_\epsilon h(1\otimes z)\otimes 1_{\mathcal{U}}. \tag{e 5.200}$$

□

The following holds in great generality and well-known. We only use the special case below. The proof in this case is easier than the general case.

**Lemma 5.4.** *Let $A$ be a unital simple AF-algebra with a unique tracial state $\tau$ and $\rho_A(K_0(A))$ is a countable dense divisible subgroup of $\mathbb{R}$. There is a unitary $u\in A$ such that*

$$\tau\circ f(u) = \int_{\mathbb{T}} f(t)dt \ \text{ for all } \ f\in C(\mathbb{T}), \tag{e 5.201}$$

*where the integral is on the standard Lesbegue measure.*

**Lemma 5.5.** *Let $A$ be a unital simple AF-algebra with a tracial state $t$ and let $v \in U(A)$. Suppose that $u \in \mathcal{U}$ is a unitary such that*

$$\tau \circ f(u) = \int_{\mathbb{T}} f(t)dt \ \text{ for all } \ f \in C(\mathbb{T}). \tag{e 5.202}$$

*Suppose $C$ is a unital separable $C^*$-algebra and $h : C \to A$ is a unital monomorphism. Define a linear map $\varphi : C \otimes C(\mathbb{T}) \to A \otimes \mathcal{U}$ by $\varphi(c \otimes f) = h(c)f(v \otimes u)$ for all $c \in C$ and $f \in C(\mathbb{T})$.*

*Then*

$$(t \otimes \tau)(\varphi(c \otimes f)) = t(h(c)) \cdot \int_{\mathbb{T}} f(t)dt \ \textit{ for all } c \in C \ \text{ and } \ f \in C(\mathbb{T}), \tag{e 5.203}$$

*where $\tau$ is the unique tracial state on $\mathcal{U}$.*

*Proof.* Let $\epsilon > 0$ and $\mathcal{F} \subset C(\mathbb{T})$ be a finite subset. We assume that $z \in \mathcal{F}$, where $z$ denotes the identity function on the unit circle.

By [31], for any $\delta > 0$, there are (for some large $n$), mutually orthogonal projections $p_1, p_2, ..., p_n$ in $A$ and mutually orthogonal projections $e_1, e_2, ..., e_n$ in $\mathcal{U}$ with $\sum_{i=1}^n p_i = 1_A$ and $\sum_{i=1}^n e_i = 1_{\mathcal{U}}$ such that

$$f(v) \approx_{\delta/4} \sum_{k=1}^n f(\omega_k)p_k \ \text{ and} \tag{e 5.204}$$

$$f(u) \approx_{\delta/4} \sum_{k=1}^n f(\omega_k)e_k \tag{e 5.205}$$

for all $f \in \mathcal{F}$, where $\omega_k = e^{\frac{2k\pi i}{n}}$ and $\tau(e_k) = 1/n$, $k = 1, 2, ..., n$. We have that

$$(\sum_{k=1}^n \omega_k p_k) \otimes (\sum_{k=1}^n \omega_k e_k) = \sum_{k=1}^n \omega_k (\sum_{j+i=k(\mathrm{mod}\, n)} p_j \otimes e_i). \tag{e 5.206}$$

Put $q_k = \sum_{j+i=k(\mathrm{mod}\, n)} p_j \otimes e_i$. Then

$$(t \otimes \tau)(q_k) = \sum_{j+i=k(\mathrm{mod}\, n)} t(p_j)\tau(e_i) \tag{e 5.207}$$

$$= \sum_{j+i=k(\mathrm{mod}\, n)} t(p_j)(1/n) = 1/n. \tag{e 5.208}$$

It follows immediately that, with sufficiently small $\delta$ (or sufficiently large $n$),

$$\tau(\varphi(1 \otimes f)) \approx_\epsilon \tau(f(u)) \ \text{ for all } \ f \in \mathcal{F}. \tag{e 5.209}$$

Furthermore, assuming $\|c\| \le 1$,

$$(t \otimes \tau)(\varphi(c \otimes f)) \approx_\epsilon (t \otimes \tau)(\sum_{k=1}^n f(\omega_k)(h(c)q_k) \tag{e 5.210}$$

$$= \sum_{k=1}^n f(\omega_k)(t \otimes \tau)((h(c)q_k) \tag{e 5.211}$$

$$= \sum_{k=1}^n f(\omega_k)(1/n)t(h(c)\sum_{j=1}^n p_j) \tag{e 5.212}$$

$$= \sum_{k=1}^n f(\omega_k)(1/n)t(h(c)) \tag{e 5.213}$$

$$\approx_\epsilon t(h(c))\tau(f(u)). \tag{e 5.214}$$

for all $f \in \mathcal{F}$. Let $\epsilon \to 0$, we conclude that

$$(t \otimes \tau)(\varphi(c \otimes f)) = t(c)\tau(f(u)) \text{ for all } c \in C \text{ and } f \in C(\mathbb{T}). \tag{e 5.215}$$

$\square$

**Lemma 5.6.** *Let $C$ be a unital AH-algebra and let $\alpha \in Aut(A)$ be an automorphism and let $A \cong A \otimes \mathcal{U}$ be a unital simple AF-algebra with a unique tracial state $\tau$ and $\rho_A(K_0(A)) = K_0(A)$. Let $\varphi : C \rtimes_\alpha \mathbb{Z} \to A$ be a unital monomorphism.*

*For any $\epsilon > 0$ and a finite subset $\mathcal{F} \subset C \rtimes_\alpha \mathbb{Z}$, there exists $\delta > 0$, a finite subset $\mathcal{G} \subset C \rtimes_\alpha \mathbb{Z}$ and a finite subset $\mathcal{P} \subset K_1(C \rtimes_\alpha \mathbb{Z})$ satisfying the following:*

*Suppose that there is a unitary $v \in U(A)$ such that*

$$\|[\varphi(a), v]\| < \delta \textit{ for all } a \in \mathcal{G} \text{ and } \mathrm{bott}_1(\varphi, v)|_{\mathcal{P}} = 0. \tag{e 5.216}$$

*Then there exists a continuous path of unitaries $\{v_t : t \in [0,1]\} \subset U(A \otimes \mathcal{U})$ such that*

$$v_0 = v \otimes 1_{\mathcal{U}}, \;\; v_1 = 1_{A \otimes \mathcal{U}}, \;\; \|[\psi(a), v_t]\| < \epsilon \tag{e 5.217}$$

*for all $a \in \mathcal{F}$ and $t \in [0,1]$, and*

$$\mathrm{Length}(\{v_t\}) \le 2\pi + \min\{\epsilon, 1\}, \tag{e 5.218}$$

*where $\psi : C \rtimes_\alpha \mathbb{Z} \to A \otimes \mathcal{U}$ defined by $\psi(a) = a \otimes 1$ for $a \in C$ and $\psi(u_\alpha) = \varphi(u_\alpha) \otimes \imath(u_\sigma)$.*

*Proof.* To simplify notation, without loss of generality, it suffices to prove the theorem with $v_t \in A \otimes \mathcal{U} \otimes \mathcal{U}$ and $\psi(u_\alpha) = \varphi(u_\alpha) \otimes 1_{\mathcal{U}} \otimes \imath(u_\sigma)$.

Let $u_0 \in \mathcal{U}$ be a unitary with

$$\tau_1(f(u_0)) = \int_{\mathbb{T}} f(t)dt \text{ for all } f \in C(\mathbb{T}), \tag{e 5.219}$$

where $\tau_1$ is the unique tracial state on $\mathcal{U}$ (by 5.4).

It follows from [31] (see 4.4.1 of [35]) that there is a continuous path of unitaries $\{V(t) : t \in [0,1]\}$ in $1 \otimes \mathcal{U}$ such that

$$V(0) = 1, \;\; V(1) = 1 \otimes u_0 \text{ and } \mathrm{Length}(\{V(t)\}) \le \pi + \epsilon/4. \tag{e 5.220}$$

Let $\epsilon > 0$ and $\mathcal{F} \subset C \rtimes_\alpha \mathbb{Z}$ be a finite subset. Without loss of generality, we may assume that $\mathcal{F} = \mathcal{F}_0 \cup \{u_\alpha\}$ for some finite subset $\mathcal{F}_0 \subset C$. Define $C_1 = C \otimes C(\mathbb{T})$. Define $\alpha' \in Aut(C_1)$ by $\alpha'(c \otimes 1) = \alpha(c) \otimes 1$ and $\alpha'(1 \otimes f) = 1 \otimes f$ for all $f \in C(\mathbb{T})$. In other words, $C_1 \rtimes_{\alpha'} \mathbb{Z} \cong (C \rtimes_\alpha \mathbb{Z}) \otimes C(\mathbb{T})$. Define $\mathcal{F}' = \mathcal{F}_0 \cup \{z\}$.

Define $h : (C \rtimes_\alpha \mathbb{Z}) \otimes C(\mathbb{T}) \to A \otimes \mathcal{U}$ by $h(a) = \varphi(a) \otimes 1_{\mathcal{U}}$ for all $a \in C \rtimes_\alpha \mathbb{Z}$ and $h(1 \otimes z) = 1 \otimes u_0$. Put $\bar{v} = v \otimes u_0$.

Let $\delta_1 > 0$ (in place of $\delta$), $\eta_1 > 0$ (in place of $\eta$), $\mathcal{G}_1 \subset C \rtimes_\alpha \mathbb{Z}$ (in place of $\mathcal{G}$) be a finite subset, $\bar{\mathcal{G}} \subset C \otimes C(\mathbb{T})$ be a finite subset and $\mathcal{P} \subset K_1(C \rtimes_\alpha \mathbb{Z})$ be a finite subset required by 5.3 for $\sin(\epsilon/4)$ and $\mathcal{F}$ above. Without loss of generality, we may assume that $\mathcal{G}_1 = \mathcal{G}_1' \cup \{u_\alpha\}$ for some finite subset $\mathcal{G}_1' \subset C$, and we may assume that $\bar{\mathcal{G}} = \mathcal{G}_2' \otimes \mathcal{G}_2''$, where $1_C \in \mathcal{G}_2' \subset C$ is a finite subset and $\{1_{C(\mathbb{T})}, z\} \subset \mathcal{G}_2'' \subset C(\mathbb{T})$ is a finite subset.

By 2.14, there is $0 < \delta < \delta_1$ and a finite subset $\mathcal{G}_3 \supset \mathcal{G}_1'$ so that there exits a unital contractive completely positive linear map $L : (C \rtimes_\alpha \mathbb{Z}) \otimes C(\mathbb{T}) \to A \otimes \mathcal{U}$ such that

$$L(c \otimes f) \approx_{\delta_1/2} \varphi(c)f(v') \text{ for all } c \in \mathcal{G}_1 \cup \mathcal{G}_2' \text{ and } f \in \mathcal{G}_2. \tag{e 5.221}$$

for any unitary $v'$ so that $\|[\varphi(a),\, v']\| < \delta$ for all $a \in \mathcal{G}_3$. Without loss of generality, we may also assume that

$$[L]|_{\mathcal{Q}} = [L']|_{\mathcal{Q}} \tag{e 5.222}$$

provided that $L, L'$ are both $\delta_1$-$\mathcal{G}_1$-multiplicative and $L \approx_{\delta_1} L'$ on $\mathcal{G}_1$.

Let $\tau_0$ be the tracial state on $C_1$ defined by

$$\tau_0(c \otimes f) = \tau(\varphi(c))\tau_1(f(u_0)) \ \text{ for all } \ c \in C \ \text{ and } \ f \in C(\mathbb{T}). \tag{e 5.223}$$

Let $\mathcal{G} = \mathcal{G}_1 \cup \mathcal{G}_2'$. Suppose that (e 5.216) holds for the above $\delta$, $\mathcal{G}$ and $\mathcal{P}$.

Then, we have that

$$\|[h(a),\, \bar{v}]\| < \delta_1 \ \text{ for all } \ a \in \mathcal{G}_1, \ \ \mathrm{bott}_1(h|_{C \rtimes_\alpha \mathbb{Z}},\, \bar{v})|_{\mathcal{P}} = 0. \tag{e 5.224}$$

Define $L_0 : C \otimes C(\mathbb{T}) \to A \otimes \mathcal{U}$ by $L_0(c \otimes f) = h(c)f(\bar{v})$ for all $c \in C$ and $f \in C(\mathbb{T})$. It follows from 5.5 that

$$\tau \circ L_0 = \tau_0, \tag{e 5.225}$$

where $\tau$ is regarded as the tracial state on $A \cong A \otimes \mathcal{U}$. There exists a contractive completely positive linear map $L : C \otimes C(\mathbb{T}) \to A \otimes \mathcal{U}$ such that (e 5.221) holds In particular,

$$\tau \circ L \approx_\eta \tau \circ h|_{C_1} \ \text{ on } \ \bar{\mathcal{G}}. \tag{e 5.226}$$

It follows from 5.3 that there exists a unitary $W \in U(A \otimes \mathcal{U} \otimes \mathcal{U})$ such that

$$\|[\psi(a),\, W]\| < \sin(\epsilon/4) \ \text{ for all } \ a \in \mathcal{F} \ \text{ and } \ W^*(\bar{v} \otimes 1_{\mathcal{U}})W \approx_{\sin(\epsilon/4)} h(1 \otimes z) \otimes 1_{\mathcal{U}}, \tag{e 5.227}$$

where $\psi : C \rtimes_\alpha \mathbb{Z} \to A \otimes \mathcal{U} \otimes \mathcal{U}$ defined by $\psi(c) = h(c) \otimes 1_{\mathcal{U}}$ for all $c \in C$ and $\psi(u_\alpha) = h(u_\alpha) \otimes \imath(u_\alpha)$.

In particular,

$$W(1 \otimes u_0 \otimes 1_{\mathcal{U}})W^* \approx_{\sin(\epsilon/4)} v \otimes u_0 \otimes 1_{\mathcal{U}}. \tag{e 5.228}$$

By (e 5.228), there exists a continuous path $\lambda : [1/4, 1/2] \to U(A \otimes \mathcal{U} \otimes \mathcal{U})$ such that

$$\begin{aligned} \lambda(1/4) &= v \otimes u_0 \otimes 1_{\mathcal{U}}, && \text{(e 5.229)}\\ \lambda(1/2) &= W(1 \otimes u_0 \otimes 1_{\mathcal{U}})W^* && \text{(e 5.230)}\\ \text{and } \ \mathrm{Length}(\{\lambda(t)\}) &< \epsilon/4 && \text{(e 5.231)} \end{aligned}$$

Moreover,

$$\|[\psi(a), \lambda(t)]\| < \delta + \frac{\epsilon}{4} < \epsilon \ \text{ for all } \ a \in \mathcal{F} \ \text{ and } \quad \text{for all } \ t \in [1/4, 1/2]. \tag{e 5.232}$$

Now define

$$v_t = \begin{cases} (v \otimes 1_{\mathcal{U}})V(4t) \otimes 1_{\mathcal{U}}, & \text{if } t \in [0, 1/4);\\ \lambda(t), & \text{if } t \in [1/4, 1/2) \text{ and}\\ W((V(2-2t)) \otimes 1_{\mathcal{U}})W^*, & \text{if } t \in [1/2, 1]. \end{cases}$$

Then

$$v_0 = v \otimes 1_{\mathcal{U}} \otimes 1_{\mathcal{U}}, \ \ v_1 = 1 \ \text{ and } \ \mathrm{Length}(\{v_t\}) < \pi + \frac{\epsilon}{4} + \frac{\epsilon}{4} + \pi + \frac{\epsilon}{4} < 2\pi + \epsilon. \tag{e 5.233}$$

For $t \in [0, 1/4)$, we estimate that

$$\|[\psi(a), v_t]\| < \delta < \epsilon \text{ for all } a \in \mathcal{F}. \tag{e 5.234}$$

By (e 5.227), for all $c \in \mathcal{F}_0$ and $t \in [1/2, 1]$,

$$\begin{aligned} \|[\psi(c), v_t]\| &= \|[W^*(\varphi(c) \otimes 1_{\mathcal{U}} \otimes 1_{\mathcal{U}})W, V(2-2t) \otimes 1_{\mathcal{U}}]\| \\ &\approx_{\epsilon/4} \|[\varphi(c) \otimes 1_{\mathcal{U}} \otimes 1_{\mathcal{U}}, V(2-2t) \otimes 1_{\mathcal{U}}]\| = 0. \end{aligned} \tag{e 5.235}$$

Note again

$$\|[\psi(u_\alpha), V(2(t-1/2) \otimes 1_{\mathcal{U}}]\| = 0. \tag{e 5.236}$$

Combining (e 5.227) and (e 5.236), for $t \in [1/2, 1]$,

$$\|[\psi(u_\alpha), v_t]\| = \|[W^*(\varphi(u_\alpha) \otimes 1_{\mathcal{U}} \otimes \imath(u_\sigma))W, V(2-2t) \otimes 1_{\mathcal{U}}]\| \tag{e 5.237}$$

$$\approx_{\epsilon/4} \|[\varphi(u_\alpha) \otimes 1_{\mathcal{U}} \otimes \imath(u_\sigma), V(2-2t) \otimes 1_{\mathcal{U}}]\| = 0. \tag{e 5.238}$$

It follows that

$$\|[\psi(c), v_t]\| < \epsilon \text{ for all } c \in \mathcal{F} \text{ and } \text{ for all } t \in [0, 1]. \tag{e 5.239}$$

□

## 6 Asymptotic unitary equivalence

**Theorem 6.1.** *Let $C$ be a unital AH-algebra and let $\alpha \in Aut(C)$ be an automorphism. Let $A \cong A \otimes \mathcal{U}$ be a unital simple AF-algebra with a unique tracial state $\tau$ and with $K_0(A) = \rho_A(K_0(A))$. Suppose that $\varphi_1, \varphi_2 : C \rtimes_\alpha \mathbb{Z} \to A$ are two unital monomorphisms such that*

$$\tau \circ \varphi_1 = \tau \circ \varphi_2 \text{ and } \tilde{\eta}(\varphi_1, \varphi_2) = 0. \tag{e 6.240}$$

*Then, there exists a continuous path of unitaries $\{w_t : t \in [0, \infty)\} \subset U(B \otimes \mathcal{U} \otimes \mathcal{U})$ such that*

$$\lim_{t \to \infty} \mathrm{ad}\, w_t \circ \psi_1''(a) = \psi_2''(a) \textit{ for all } a \in C \rtimes_\alpha \mathbb{Z}, \tag{e 6.241}$$

*where $B = B(\psi_2, A \otimes \mathcal{U})$ and where $\psi_i : C \rtimes_\alpha \mathbb{Z} \to A \otimes \mathcal{U}$ is defined by $\psi_i(c) = \varphi_i(c) \otimes 1_{\mathcal{U}}$ for all $a \in C$ and $\psi_i(u_\alpha) = \varphi_i(u_\alpha) \otimes \imath(u_\sigma)$, and $\psi_i'' : C \rtimes_\alpha \mathbb{Z} \to B \otimes \mathcal{U} \otimes \mathcal{U}$ is defined by $\psi_i''(c) = \varphi(c) \otimes 1_{\mathcal{U}} \otimes 1_{\mathcal{U}}$ for $c \in C$ and $\psi_i''(u_\alpha) = \psi(u_\alpha) \otimes \imath(u_\alpha)$, $i = 1, 2$.*

*Proof.* Since $\ker \rho_A = \{0\}$, (e 6.240) implies that

$$(\varphi_1)_{*0} = (\varphi_2)_{*0}. \tag{e 6.242}$$

Since $K_1(A) = \{0\}$ and $K_0(A)$ is torsion free and divisible, (e 6.242) implies that

$$[\varphi_1] = [\varphi_2] \text{ in } KK(C \rtimes_\alpha \mathbb{Z}, A). \tag{e 6.243}$$

Therefore, since $\tilde{\eta}_{\varphi_1, \varphi_2} = 0$, there exists $\theta : K_1(C \rtimes_\alpha \mathbb{Z}) \to K_1(M_{\varphi_1, \varphi_2})$ such that

$$(\pi_0)_{*1} \circ \theta = (\mathrm{id}_{C \rtimes_\alpha \mathbb{Z}})_{*1}. \tag{e 6.244}$$

Moreover,

$$R_{\varphi_1, \varphi_2}(\theta(K_1(C \rtimes_\alpha \mathbb{Z}))) = 0. \tag{e 6.245}$$

Denote by $j_0 : C \to C \rtimes_\alpha \mathbb{Z}$ the natural embedding. Then one has

$$R_{\varphi_1,\varphi_2}(\theta \circ (j_0)_{*1}(K_1(C))) = 0. \tag{e 6.246}$$

Therefore

$$\tilde{\eta}_{\varphi_1 \circ j_0, \varphi_2 \circ j_0} = 0. \tag{e 6.247}$$

It follows from 3.6 that there exists a sequence of unitaries $\{v_n\} \subset A \otimes \mathcal{U}$ such that

$$\lim_{n\to\infty} \mathrm{ad}\, v_n \circ \psi_1(a) = \psi_2(a) \ \text{ for all } \ a \in C \rtimes_\alpha \mathbb{Z}, \tag{e 6.248}$$

where $\psi_i : C \rtimes_\alpha \mathbb{Z} \to A \otimes \mathcal{U}$ defined by $\psi_i(c) = \varphi_i(c) \otimes 1_{\mathcal{U}}$ for all $c \in C$ and $\psi_i(u_\alpha) = \varphi_i(u_\alpha) \otimes \imath(u_\alpha)$.

Let $B = B(\psi_2, A \otimes \mathcal{U})$ be as in 4.3.

To simplify notation, without loss of generality, in what follows we will use $\tau$ for the unique tracial state on $A$, $A \otimes \mathcal{U}$ $(\cong A)$, $B \otimes \mathcal{U}$ and $B \otimes \mathcal{U} \otimes \mathcal{U} \cong B \otimes \mathcal{U}$.

Let $\{\mathcal{F}_n\} \subset C \rtimes_\alpha \mathbb{Z}$ be an increasing sequence of finite subsets of the unit ball of $C \rtimes_\alpha \mathbb{Z}$ whose union is dense in the unit ball. Let $\delta_n > 0$, $\mathcal{G}_n \subset C \rtimes_\alpha \mathbb{Z}$ be a finite subset and $\mathcal{P}_n \subset K_1(C \rtimes_\alpha \mathbb{Z})$ be a finite subset required by 5.6 for $\epsilon_n = 1/2^{n+2}$ and $\mathcal{F}_n$. We may assume that $\mathrm{bott}_1(h', v')|_{\mathcal{P}_n}$ is well defined for any unital homomorphism $h'$ and any unitary $v'$, whenever

$$\|[h'(c), v']\| < \delta_n \ \text{ for all } \ c \in \mathcal{G}_n \tag{e 6.249}$$

for $n = 1, 2, ....$ We also assume that

$$\mathrm{bott}_1(h', v')|_{\mathcal{P}_n} = \mathrm{bott}_1(h', v'')|_{\mathcal{P}_n} \tag{e 6.250}$$

provided that $\|v' - v''\| < \delta_n$.

We may assume that $\delta_n < 1/2^{n+1}$ and $\mathcal{F}_n \subset \mathcal{G}_n$, $n = 1, 2, ....$ We may write $\mathcal{P}_n = \{x_1, x_2, ..., x_{r(n)}\}$, $n = 1, 2, ....$

Let $\delta_n' > 0$ and $\{y_1, y_2, ..., y_{m(n)}\} \subset K_1(C \rtimes_\alpha \mathbb{Z})$ be a finite subset required by 4.4 for $\delta_n/2$, $\mathcal{G}_n$ and $\mathcal{P}_n$. Put $\mathcal{Q}_n = \{y_1, y_2, ..., y_{m(n)}\}$. We may assume that $m(n+1) \geq m(n)$. To simplify notation, we may assume that $x_j = y_j$, $j = 1, 2, ..., r(n)$ and $r(n) \leq m(n)$, $n = 1, 2, ....$

There is an integer $l(n) \geq 1$ and unitaries $z_1, z_2, ..., z_{m(n)} \in M_{l(n)}(C \rtimes_\alpha \mathbb{Z})$ such that $[z_j] = y_j$, $j = 1, 2, ..., m(n)$.

Let $\eta_n > 0$ be such that $\eta_n < \min\{\delta_n/2l(n)^2, (1/4)\sin(\delta_n'/2l(n)^2)\}$ and $\eta_{n+1} < \eta_n/2$, $n = 1, 2, ....$ We may also assume that, by passing to a subsequence, if necessary

$$\mathrm{ad}\, v_n \circ \psi_1 \approx_{\eta_n/2} \psi_2 \ \text{ on } \ \mathcal{G}_n \tag{e 6.251}$$

Note that

$$\|[\psi_1(a), v_n v_{n+1}^*]\| < \eta_n \ \text{ for all } \ a \in \mathcal{G}_n \tag{e 6.252}$$

So $\mathrm{bott}_1(\varphi_1, v_n v_{n+1}^*)|_{\mathcal{Q}_n}$ is well-defined.

Put

$$\bar{v}_n = \mathrm{diag}(\overbrace{v_n, v_n, ..., v_n}^{l(n)}).$$

We will continue to use $\varphi_i$ for $\varphi_i \otimes \mathrm{id}_{M_{l(n)}}$ and use $\psi_i$ for $\psi_i \otimes \mathrm{id}_{M_{l(n)}}$, $i = 1, 2$, respectively. Thus, we may assume that, for $j \leq m(n)$,

$$\|\psi_2(z_j) \mathrm{ad}\, \bar{v}_n \circ \psi_1(z_j^*) - 1\| < (1/4)\sin(\delta_n'/2) \tag{e 6.253}$$

Put

$$h_{j,n} = \log(\frac{1}{2\pi i}\psi_2(z_j)\mathrm{ad}\,\bar{v}_n \circ \psi_1(z_j^*)),\ \ j=1,2,...,m(n),\ n=1,2,.... \tag{e 6.254}$$

Then

$$\tau(h_{j,n}) < \delta_n/2, \tag{e 6.255}$$

By (e 6.246), $R_{\psi_1,\psi_2}(K_1(M_{\psi_1,\psi_2})) \subset \rho_{A\otimes\mathcal{U}}(K_0(A\otimes\mathcal{U}))$. It follows from 3.5 of [49] that

$$\widehat{h_{j,n}}(\tau) = \tau(h_{j,n}) \in \rho_{A\otimes\mathcal{U}}(K_0(A\otimes\mathcal{U})),\ \ j=1,2,...,m(n), n=1,2,.... \tag{e 6.256}$$

Since $K_0(A\otimes\mathcal{U})$ is divisible, there exists a homomorphism $\beta_n : K_1(C\rtimes_\alpha\mathbb{Z}) \to K_0(A\otimes\mathcal{U})$ (see also 6.1, 6.2 and 6.3 of [48]) such that

$$\tau(\beta_n(y_j)) = \widehat{h_{j,n}}(\tau) = \tau(h_{j,n}), j=1,2,...,m(n), n=1,2,.... \tag{e 6.257}$$

It follows from 4.4 that there exists a unitary $U_n \in B = B(\psi_2, A\otimes\mathcal{U})$ such that

$$\|[\psi_2'(a),\, U_n]\| < \delta_n' \ \text{ for all } \ a\in\mathcal{G}_n \ \text{ and } \ \mathrm{bott}_1(\psi_2',\, U_n)|_{\mathcal{P}_n} = -\beta_n'|_{\mathcal{P}_n}, \tag{e 6.258}$$

where $\beta_n' = (j_2)_{*0}\circ\beta_n$, $n=1,2,...,$ and $j_2 : A\otimes\mathcal{U} \to B$ is the embedding and where $\psi_i : C\rtimes_\alpha\mathbb{Z} \to B\otimes\mathcal{U}$ is defined by $\psi_i'(c) = \psi(c)\otimes 1_{\mathcal{U}}$ for all $c\in C\rtimes_\alpha\mathbb{Z}$ and $\psi(u_\alpha)\otimes\imath(u_\alpha)$. Note that (see 4.3) that $\tau\circ j_2(a) = \tau(a)$ for $a\in A\otimes\mathcal{U}$.

By Exel's trace formula (see Theorem 3.6 of [49]) for $j=1,2,...,r(n)$,

$$\tau(h_{j,n}) = -\mathrm{bott}_1(\psi_2',\, U_n)(x_j)(\tau) \tag{e 6.259}$$

$$= -\tau(\log(\frac{1}{2\pi i}\bar{U}_n\psi_2'(z_j)\bar{U}_n^*\psi_2'(z_j^*))), \tag{e 6.260}$$

where

$$\bar{U}_n = \mathrm{diag}(\overbrace{U_n, U_n, ..., U_n}^{l(n)}).$$

Define $u_n = v_nU_n$, $n=1,2,....$ Put

$$\bar{u}_n = \mathrm{diag}(\overbrace{u_n, u_n, ..., u_n}^{l(n)}) \ \text{ and } \ \bar{u}_{n+1}' = \mathrm{diag}(\overbrace{u_{n+1}, u_{n+1}, ..., u_{n+1}}^{l(n)}), \tag{e 6.261}$$

$n=1,2,....$ By 6.1 of [48] and (e 6.259), we compute that

$$\tau(\log(\frac{1}{2\pi i}(\psi_2'(z_j)\mathrm{ad}\,\bar{u}_n(\psi_1'(z_j^*))))) \tag{e 6.262}$$

$$= \tau(\log(\frac{1}{2\pi i}(\bar{U}_n\psi_2'(z_j^*)\bar{U}_n^*\bar{v}_n^*(\psi_1'(z_j))\bar{v}_n))) \tag{e 6.263}$$

$$= \tau(\log(\frac{1}{2\pi i}(\bar{U}_n\psi_2'(z_j)\bar{U}_n^*\psi_2'(z_j^*)\psi_2'(z_j)\bar{v}_n^*(\psi_1'(z_j^*))\bar{v}_n))) \tag{e 6.264}$$

$$= \tau(\log(\frac{1}{2\pi i}(\bar{U}_n\psi_2'(z_j)\bar{U}_n^*\psi_2'(w_j^*)))) \tag{e 6.265}$$

$$+\tau(\frac{1}{2\pi i}\log(\psi_2'(z_j)\bar{v}_n^*(\psi_1'(z_j^*))\bar{v}_n))) \tag{e 6.266}$$

$$= \mathrm{bott}_1(\psi_2', U_n))(z_j)(\tau) + \tau(h_{j,n}) = 0. \tag{e 6.267}$$

Let

$$\begin{align} b_{j,n} &= \log(\frac{1}{2\pi i}\bar{u}_n\psi_2'(z_j)\bar{u}_n^*\psi_1'(z_j^*)) \text{ and} \tag{e 6.268}\\ b_{j,n}' &= \log(\frac{1}{2\pi i}\psi_2'(z_j)\bar{u}_n^*\bar{u}_{n+1}'\psi_2'(z_j^*)(\bar{u}_{n+1}')^*\bar{u}_n)), \tag{e 6.269}\end{align}$$

$j = 1, 2, ..., n$ and $n = 1, 2, ....$ We have, by (e 6.267),

$$\begin{align} \tau(b_{j,n}) &= \tau(\log(\frac{1}{2\pi i}\bar{u}_n\psi_2'(z_j)\bar{u}_n^*\psi_1'(z_j^*))) \tag{e 6.270}\\ &= \tau(\log\frac{1}{2\pi i}\bar{u}_n^*\bar{u}_n\psi_2'(z_j)\bar{u}_n^*\psi_1'(z_j^*)\bar{u}_n) \tag{e 6.271}\\ &= \tau(\log\frac{1}{2\pi i}\psi_2'(z_j)\bar{u}_n^*\psi_1'(z_j^*)\bar{u}_n) = 0. \tag{e 6.272}\end{align}$$

$j = 1, 2, ..., r(n)$ and $n = 1, 2, ....$ Note also $\tau(b_{j,n+1}) = 0$ for $j = 1, 2, ..., r(n)$. In $M_{l(n+1)}(C \rtimes_\alpha \mathbb{Z})$, $x_j$ represented by

$$\mathrm{diag}(z_j, 1_{M_{l(n+1)-l(n)}}) \tag{e 6.273}$$

for $j = 1, 2, ..., r(n)$. Put

$$b_{j,n+1}'' = \log(\frac{1}{2\pi i}\psi_2'(z_j)(\bar{u}_{n+1}')^*\psi_1'(z_j^*)\bar{u}_{n+1}') \tag{e 6.274}$$

Therefore

$$\tau(b_{j,n+1}'') = \tau(b_{j,n+1}) = 0. \tag{e 6.275}$$

We have that

$$\bar{u}_n^* e^{2\pi i b_{j,n}'}\bar{u}_n = e^{2\pi i b_{j,n}}e^{-2\pi i b_{j,n+1}''}. \tag{e 6.276}$$

Thus, by 6.1 of [48], we compute that

$$\tau(b_{j,n}') = \tau(b_{j,n}) - \tau(b_{j,n+1}) = 0. \tag{e 6.277}$$

It follows the Exel trace formula (Theorem 3.6 of [49]) and (e 6.277) that

$$\begin{align} &\mathrm{bott}_1(\psi_2', u_n^*u_{n+1}))(x_j)(\tau) \tag{e 6.278}\\ &= -\tau(\log(\bar{u}_n^*\bar{u}_{n+1}'\psi_2(z_j^*)(\bar{u}_{n+1}')^*\bar{u}_n\psi_2(z_j)) \tag{e 6.279}\\ &= -\tau(\log(\psi_2'(z_j)\bar{u}_n^*\bar{u}_{n+1}'\psi_2'(z_j^*)(\bar{u}_{n+1}')^*\bar{u}_n)) = 0. \tag{e 6.280}\end{align}$$

By applying 5.6, we obtain a continuous path of unitaries $\{v_n(t) : t \in [0,1]\} \subset B \otimes \mathcal{U} \otimes \mathcal{U}$ such that

$$v_n(0) = u_n u_{n+1}^*, \;\; v_n(1) = 1 \;\text{ and }\; \|[\psi_2''(a),\, v_n(t)]\| < 1/2^{n+2} \;\text{ for all }\; a \in \mathcal{F}_n, \tag{e 6.281}$$

where $\psi_i'' : C \rtimes_\alpha \mathbb{Z} \to B \otimes \mathcal{U} \otimes \mathcal{U}$ is defined by $\psi_i''(c) = \psi_i'(c)$ for all $c \in C$ and $\psi_i''(u_\alpha) = \psi_i'(u_\alpha) \otimes \imath(u_\sigma)$, $i = 1, 2$ and $n = 1, 2, ....$ Define $u(t) = v_n(t - n + 1)u_{n+1}$ for $t \in [n-1, n)$, $n = 1, 2, ....$ We check that

$$\lim_{t\to\infty} \mathrm{ad}\, u(t) \circ \psi_1''(a) = \psi_2''(a) \;\text{ for all }\; a \in C \rtimes_\alpha \mathbb{Z}. \tag{e 6.282}$$

□

# 7 AF-embedding of $C\rtimes_\Lambda \mathbb{Z}^2$

**Theorem 7.1.** *Let $C$ be a unital AH-algebra and let $\Lambda: \mathbb{Z}^2\to Aut(C)$ be a homomorphism. Then $C\rtimes_\Lambda \mathbb{Z}^2$ can be embedded into a unital simple AF-algebra if and only if $C$ admits a faithful $\Lambda$-invariant tracial state.*

*Proof.* We only need to show the "if " part of the theorem. Let $\alpha_1$ and $\alpha_2$ be two generators of $\Lambda(\mathbb{Z}^2)$. Thus $\alpha_1, \alpha_2\in Aut(A)$, $\alpha_1\circ \alpha_2=\alpha_2\circ \alpha_1$ and $C\rtimes_\Lambda \mathbb{Z}^2\cong (C\rtimes_{\alpha_1}\mathbb{Z})\rtimes_{\alpha_2}\mathbb{Z}$.

Put $C_1=C\rtimes_{\alpha_1}\mathbb{Z}$. We view $\alpha_2$ as an automorphism on $C_1$ by $\alpha_2(u_{\alpha_1})=u_{\alpha_1}$.

Let $t$ be a faithful $\Lambda$-invariant tracial state on $C$. We may also view $t$ as a faithful tracial state on $C_1$ as well as on $C\rtimes_\Lambda \mathbb{Z}^2$.

It follows from (the proof) of 10.5 of [49] that there exists a unital simple AF-algebra $A\cong A\otimes \mathcal{U}$ with unique tracial state $\tau$ and with $K_0(A)=\rho_A(K_0(A))$ and a unital monomorphism $h: C_1\to A$ such that

$$\tau\circ h=t. \tag{e 7.283}$$

Put $\varphi_1=h$ and $\varphi_2=h\circ \alpha_2$. Then

$$\tau\circ \varphi_1=\tau\circ \varphi_2. \tag{e 7.284}$$

In particular, since $\mathrm{ker}\rho_A=\{0\}$, $K_1(A)=\{0\}$ and $K_0(A)$ is torsion free and divisible,

$$[\varphi_1]=[\varphi_2]\ \ \text{in}\ \ KK(C_1,A). \tag{e 7.285}$$

There exists $\theta: K_1(C_1)\to K_1(M_{\varphi_1,\varphi_2})$. Let $\mathbb{D}_0=R_{\varphi_1,\varphi_2}(\theta(K_1(C_1)))$. Let $\mathbb{D}_1$ be a countable subring of $\mathbb{R}$ containing $\rho_A(K_0(A))\cup \mathbb{D}_0$ which is also a divisible subgroup of $\mathbb{R}$. Let $A_1$ be a unital simple AF-algebra with a unique tracial state such that $K_0(A_1)=\mathbb{D}_1$. Moreover, there is a unital embedding $j: A\to A_1$ such that

$$\tau\circ j=\tau, \tag{e 7.286}$$

where we also use $\tau$ for the unique tracial state on $A_1$. To simplify notation, without loss of generality, we may assume that $A=A_1$. Therefore, with this assumption, we have

$$R_{\varphi_1,\varphi_2}(M_{\varphi_1,\varphi_2})\subset \rho_A(K_0(A))=\mathbb{D}_1. \tag{e 7.287}$$

By 10.3 of [49], this implies that

$$\tilde{\eta}_{\varphi_1,\varphi_2}=0. \tag{e 7.288}$$

It follows that 6.1 that there exists a continuous path of unitaries $\{w_t: t\in [0,\infty)\}$ of $B\otimes \mathcal{U}\otimes \mathcal{U})$ such that

$$\lim_{t\to\infty}\mathrm{ad}\, w_t\circ \psi_1''(a)=\psi_2''(a)\ \ \text{for all}\ \ a\in C_1, \tag{e 7.289}$$

where $\psi_i'': C_1\to B\otimes \mathcal{U}\otimes \mathcal{U}$ is defined by $\psi_i''(c)=\varphi(c)\otimes 1_{\mathcal{U}}\otimes 1_{\mathcal{U}}$ for all $c\in C$ and $\psi_i''(u_{\alpha_1})=\varphi_i(u_{\alpha_1})\otimes\imath(u_\sigma)\otimes\imath(u_\sigma)$, and $B=B(\psi_2, A\otimes\mathcal{U})$ and $\psi_2: C_1\to A\otimes\mathcal{U}$ is defined by $\psi_2(c)=\varphi_2(c)\otimes 1_{\mathcal{U}}$ for all $c\in C$ and $\psi_2(u_{\alpha_1})=\varphi_2(u_{\alpha_1})\otimes \imath(u_\sigma)\otimes\imath(u_\sigma)$. It follows from 5.5 of [48] that there exists a unital monomorphism $h_1: C_1\rtimes_{\alpha_2}\mathbb{Z}\to B_1$, where $B_1$ is a unital simple AF-algebra with a unique tracial state.

□

**Corollary 7.2.** *Let $C$ be a unital AH-algebra and let $\Lambda : \mathbb{Z}^2 \to Aut(C)$ be a homomorphism. Suppose that $t$ is a faithful $\Lambda$-invariant tracial state on $C$.*

*Then there exists a unital simple AF-algebra $B_1$ with a unique tracial state $\tau$ and with $K_0(B_1) = \rho_{B_1}(K_0(B_1))$ which is also a subring of $\mathbb{R}$, and there exists a unital monomorphism $h_1 : C \rtimes_\Lambda \mathbb{Z}^2 \to B_1$ such that*

$$\tau \circ h_1 = t. \tag{e 7.290}$$

*Proof.* At the end of the proof of 7.1, we apply 5.5 of [48]. The proof of 5.5 of [48] actually shows that one can have the further requirements for $B_1$, $h_1$ and $\tau$.

□

**Corollary 7.3.** *Let $X$ be a compact metric space and $\Lambda : \mathbb{Z}^2 \to Homeo(X)$ be a homomorphism. Then $C(X) \rtimes_\Lambda \mathbb{Z}^2$ can be embedded into a unital simple AF-algebra if and only if $X$ admits a strictly positive $\Lambda$-invariant probability Borel measure.*

## 8 The absorption lemma

**Definition 8.1.** Let $C$ be a unital separable amenable $C^*$-algebra and let $\Lambda_k : \mathbb{Z}^k \to Aut(C)$ be a homomorphism. Suppose that $\{\alpha_1, \alpha_2, ..., \alpha_k\}$ is the set of generators of $\Lambda(\mathbb{Z}^k)$.

Suppose that $\varphi : C \rtimes_\Lambda \mathbb{Z}^k \to A$ is a unital monomorphism. Denote by $\varphi^{(k)} : C \rtimes_{\Lambda_1} \mathbb{Z} \to A \otimes \mathcal{U}^k$ the monomorphism defined by

$$\varphi^{(k)}(c) = \varphi(c) \otimes 1_{\mathcal{U}^k} \quad \text{for all } c \in C, \tag{e 8.291}$$

$$\varphi^{(k)}(u_{\alpha_1}) = \varphi(u_{\alpha_1}) \otimes \imath(u_\sigma) \otimes 1_{\mathcal{U}^{k-1}}, \tag{e 8.292}$$

$$\varphi^{(k)}(u_{\alpha_2}) = \varphi(u_{\alpha_2}) \otimes 1_{\mathcal{U}} \otimes \imath(u_\sigma) \otimes 1_{\mathcal{U}^{k-1}}, \tag{e 8.293}$$

$$\cdots \quad \cdots \tag{e 8.294}$$

$$\varphi^{(k)}(u_{\alpha_k}) = (\varphi(u_{\alpha_k}) \otimes 1_{\mathcal{U}^{(k-1)}}) \otimes \imath(u_\sigma). \tag{e 8.295}$$

Let $\{\alpha_1, \alpha_2, ..., \alpha_k\}$ be a set of generators for $\Lambda(\mathbb{Z}^k)$. Define $\Lambda_j : \mathbb{Z}^j \to Aut(C)$ by $\{\alpha_1, \alpha_2, ..., \alpha_j\}$, $j = 1, 2, ..., k$. Denote by $\varphi_j : C \rtimes_{\Lambda_j} \mathbb{Z}^j \to A$ the homomorphism $\varphi|_{C \rtimes_{\Lambda_j} \mathbb{Z}^j}$ and $\varphi^{(j)} = \varphi_j^{(j)}$, $j = 1, 2, ..., k$.

Denote by $B_1 = B(\varphi, A)$ the AF-algebra defined in 4.3. Define $B_{j+1}(\varphi, A) = B(\varphi^{(j+1)}, B_j \otimes \mathcal{U}^j)$, $j = 1, 2, ..., k-1$.

**Lemma 8.2.** *In $\mathcal{U}$, there is a continuous and piecewise smooth path of unitaries $\{u(t) : t \in [0,1]\}$ in $\mathcal{U}$ such that*

$$u(0) = \imath(u_\sigma),\ \ u(1) = 1_{\mathcal{U}} \text{ and} \tag{e 8.296}$$

$$\int_0^1 \tau(\frac{du(t)}{dt} u(t)^*) dt = 0, \tag{e 8.297}$$

*where $\tau$ is the unique tracial state of $\mathcal{U}$.*

*Proof.* There are two self-adjoint elements $h_1, h_2 \in \mathcal{U}$ such that

$$\varphi(u_\sigma) = e^{2\pi i h_1} e^{2\pi i h_2}, \tag{e 8.298}$$

where $\varphi : \mathcal{U} \rtimes_\sigma \mathbb{Z} \to \mathcal{U}$ is defined in 2.11 (see [31]).

Define $v(t) = e^{2\pi i h_1(1-t)} e^{2\pi(ih_2(1-t)}$. Then $v(0) = \varphi(u_\sigma)$ and $v(1) = 1_{\mathcal{U}}$. Define $w(t) = v(t) \otimes v(t)^*$ for $t \in [0,1]$. Then,

$$\begin{aligned} \frac{dw(t)}{dt} w(t)^* &= \frac{d((v(t) \otimes 1_{\mathcal{U}})(1_{\mathcal{U}} \otimes v(t)^*)}{dt} ((v(t)^* \otimes 1_{\mathcal{U}})(1_{\mathcal{U}} \otimes v(t))) && \text{(e 8.299)} \\ &= (-ih_1) \otimes 1_{\mathcal{U}} + (-ie^{2\pi i h_1(1-t)} h_2 e^{-2\pi i h_1(1-t)}) \otimes 1_{\mathcal{U}} && \text{(e 8.300)} \\ &+ 1_{\mathcal{U}} \otimes (ih_1) \otimes 1_{\mathcal{U}} + (ie^{2\pi i h_1(t-1)} h_2 e^{2\pi i h_1(1-t)}) \otimes 1_{\mathcal{U}} && \text{(e 8.301)} \end{aligned}$$

Therefore

$$\tau(\frac{dw(t)}{dt} w(t)^*) = 0. \qquad \text{(e 8.302)}$$

Define $u(t) = s(w(t))$ for $t \in [0,1]$, where $s$ is defined in 2.11. Note that $u(0) = s(w(0)) = s(\varphi(u_\sigma) \otimes \varphi(u_\sigma)^*) = \imath(u_\sigma)$ and $u(1) = 1_{\mathcal{U}}$. Moreover,

$$\begin{aligned} \tau(\frac{du(t)}{dt} u(t)^*) &= \tau(s(\frac{dw(t)}{dt} w(t)^*)) && \text{(e 8.303)} \\ &= \tau(\frac{dw(t)}{dt} w(t)^*) = 0 && \text{(e 8.304)} \end{aligned}$$

for all $t \in [0,1]$.

□

**Lemma 8.3.** *There exists a continuous path of unitaries* $\{w(t) : t \in [0,\infty)\}$ *of* $\mathcal{U} \otimes \mathcal{U}$ *such that*

$$\lim_{t\to\infty} w(t)^*(\imath(u_\sigma) \otimes \imath(u_\sigma)) w(t) = \imath(u_\sigma) \otimes 1_{\mathcal{U}}. \qquad \text{(e 8.305)}$$

*Proof.* Let $\tau$ be the unique tracial state of $\mathcal{U}$. Using the standard conditional expectation, one computes that

$$\tau(\imath(u_\sigma)) = 0. \qquad \text{(e 8.306)}$$

Let $f = \sum_{i=1}^n \lambda_i u_\sigma^i$ be a polynomial of $u_\sigma$, where $\lambda_i \in \mathbb{C}$. Then

$$\tau(\imath(f)) = \lambda_0. \qquad \text{(e 8.307)}$$

Similarly

$$\tau(\sum_{i=1}^n \lambda_i (\imath(u_\sigma) \otimes \imath(u_\sigma))^i) = \lambda_0. \qquad \text{(e 8.308)}$$

It follows that

$$\tau(f(\imath(u_\sigma))) = \tau(f(\imath(u_\sigma) \otimes \imath(u_\sigma))) \qquad \text{(e 8.309)}$$

for all $f \in C(\mathbb{T})$, where we also use $\tau$ for the unique tracial state of $\mathcal{U} \otimes \mathcal{U}$.

Define $h_1, h_2 : C(\mathbb{T}) \to \mathcal{U} \otimes \mathcal{U}$ by $h_1(f) = f(\imath(u_\sigma) \otimes \imath(u_\sigma))$ and $h_2(f) = f(\imath(u_\sigma)) \otimes 1_{\mathcal{U}}$ for all $f \in C(\mathbb{T})$. Since $K_1(\mathcal{U}) = \{0\}$, by (e 8.309), we have

$$[h_1] = [h_2] \text{ in } KK(C(\mathbb{T}), \mathcal{U}) \text{ and } \tau \circ h_1 = \tau \circ h_2. \qquad \text{(e 8.310)}$$

Define $\theta : C(\mathbb{T}) \to M_{h_1,h_2}$ by

$$\theta(f)(t) = \imath(u_\sigma) \otimes u(t) \text{ for all } t \in [0,1], \qquad \text{(e 8.311)}$$

where $u(t)$ is a continuous piecewise smooth path of unitaries in $\mathcal{U}$ such that

$$u(0) = \imath(u_\sigma), \;\; u(1) = 1_{\mathcal{U}} \;\text{ and }\; \int_0^1 \tau(\frac{du(t)}{dt}u(t)^*)dt = 0, \tag{e 8.312}$$

given by 8.2. Then $\tilde{\eta}_{h_1,h_2} = 0$. By applying the main theorem of [49], there exists a continuous path of unitaries $\{w(t) : t \in [0, \infty)\} \subset \mathcal{U} \otimes \mathcal{U}$ such that

$$\lim_{t\to\infty} w(t)^*(\imath(u_\sigma) \otimes \imath(u_\sigma))w(t) = \imath(u_\sigma). \tag{e 8.313}$$

□

The proof of the following is similar to but easier than that of 8.3.

**Lemma 8.4.** *There exists a continuous path of unitaries* $\{u(t) : t \in [0, \infty)\} \subset \mathcal{U} \otimes \mathcal{U}$ *such that*

$$\lim_{t\to\infty} u(t)^*(\imath(u_\alpha) \otimes 1_{\mathcal{U}})u_t = 1_{\mathcal{U}} \otimes \imath(u_\alpha). \tag{e 8.314}$$

**Lemma 8.5.** *Let $C$ be a unital separable amenable $C^*$-algebra, let $\Lambda : \mathbb{Z}^k \to Aut(A)$ be a homomorphism such that $\{\alpha_1, \alpha_2, ..., \alpha_k\}$ forms a set of generators for $\Lambda(\mathbb{Z}^k)$. Let $\varphi : C \rtimes_\Lambda \mathbb{Z}^k \to A$ be a monomorphism. Then there exists a continuous path of unitaries $\{w(t) : t \in [0, \infty)\}$ of $A \otimes \mathcal{U}^{2k}$ such that*

$$\lim_{t\to\infty} \mathrm{ad}\, w(t) \circ (\varphi^{(k)})^{(k)}(a) = \varphi^{(k)}(a) \otimes 1_{\mathcal{U}^k} \;\text{ for all }\; a \in C. \tag{e 8.315}$$

*Proof.* Define $\psi : C \rtimes_\Lambda \mathbb{Z}^k \to A \otimes \mathcal{U}^{2k}$ as follows:

$$\begin{aligned}
\psi(c) &= \varphi(c) \otimes \overbrace{1_{\mathcal{U}} \oplus 1_{\mathcal{U}} \oplus \cdots \oplus 1_{\mathcal{U}}}^{2k} \qquad \text{for all } c \in C, && \text{(e 8.316)}\\
\psi(u_{\alpha_1}) &= \varphi(u_{\alpha_1}) \otimes \imath(u_\sigma) \otimes \imath(u_\sigma) \otimes 1_{\mathcal{U}^{2k-2}}, && \text{(e 8.317)}\\
\psi(u_{\alpha_2}) &= \varphi(u_{\alpha_2}) \otimes 1_{\mathcal{U}^2} \oplus \imath(u_\sigma) \otimes \imath(u_\sigma) \otimes 1_{\mathcal{U}^{2k-4}}, && \text{(e 8.318)}\\
\cdots & \;\;\cdots && \text{(e 8.319)}\\
\psi(u_{\alpha_k}) &= \varphi(u_{\alpha_k}) \oplus 1_{\mathcal{U}^{2k-2}} \oplus \imath(u_\sigma) \otimes \imath(u_\sigma). && \text{(e 8.320)}
\end{aligned}$$

It is clear, by 8.4, that there exists a continuous path of unitaries $\{u(t) : t \in [0, \infty)\} \subset \mathcal{U}^{2k}$ such that

$$\lim_{t\to\infty} (1 \otimes u(t))^*(\varphi^{(k)})^{(k)}(a)(1 \otimes u(t)) = \psi(a) \;\text{ for all }\; a \in C. \tag{e 8.321}$$

Therefore, to complete the proof, it suffices to show that there exists a continuous path of unitaries $\{w(t) : t \in [0, \infty)\}$ of $A \otimes \mathcal{U}^{2k}$ such that

$$\lim_{t\to\infty} \mathrm{ad}\, w(t) \circ \psi(a) = \varphi^{(k)}(a) \otimes 1_{\mathcal{U}^k} \;\text{ for all }\; a \in C. \tag{e 8.322}$$

But this clearly follows from 8.3.

□

**Corollary 8.6.** *Let $C$ be a unital AH-algebra and let $\alpha \in Aut(C)$ be an automorphism. Let $A \cong A \otimes \mathcal{U}$ be a unital simple AF-algebra with a unique tracial state $\tau$ and with $K_0(A) = \rho_A(K_0(A))$. Suppose that $\varphi_1, \varphi_2 : C \rtimes_\alpha \mathbb{Z} \to A$ are two unital monomorphisms such that*

$$\tau \circ \varphi_1 = \tau \circ \varphi_2 \;\text{ and }\; \tilde{\eta}(\varphi_1, \varphi_2) = 0. \tag{e 8.323}$$

*Then, there exists a continuous path of unitaries* $\{w_t : t \in [0, \infty)\} \subset U(B \otimes \mathcal{U})$ *such that*

$$\lim_{t\to\infty} w_t^*(\psi_1(a) \otimes 1_{\mathcal{U}})w_t = \psi_2(a) \otimes 1_{\mathcal{U}} \textit{ for all } a \in C \rtimes_\alpha \mathbb{Z}, \tag{e 8.324}$$

*where* $B = B(\varphi_2^{(1)}, A \otimes \mathcal{U})$ *and with* $1_A = 1_B$*, and* $\psi_i = j_2 \circ \varphi_i^{(1)}$ *(where* $j_2 : A \otimes \mathcal{U} \to B$ *is defined in* 4.3*),* $i = 1, 2$.

*Proof.* This follows immediately from 6.1 and 8.5. □

**Lemma 8.7.** *Let* $B_1 \cong B_1 \otimes \mathcal{U}$ *and* $B_2 \cong B_2 \otimes \mathcal{U}$ *be two unital simple AF-algebras with unique tracial state and with* $\rho_{B_i}(K_0(B_i)) = K_0(B_i)$ *which is also a subring of* $\mathbb{R}$ *(*$i = 1, 2$*). Suppose also that* $B_1 \otimes \mathcal{U}^k \subset B_2$*,* $1_{B_1 \otimes \mathcal{U}^k} = 1_{B_2}$ *for some* $k \geq 1$ *and suppose that* $j$ *is the embedding from* $B_1 \otimes \mathcal{U}^k$ *to* $B_2$*. Let* $j_0 : B_1 \to B_2$ *be a unital embedding such that* $(j_0)_{*0} = \mathrm{j}_{*0}$*. Then there is a continuous path of unitaries* $\{u_t : t \in [0, \infty)\} \subset B_2 \otimes \mathcal{U}^k$ *such that*

$$\lim_{t\to\infty} u_t^*(a \otimes b)u_t = j_0(a) \otimes b \tag{e 8.325}$$

*for all* $a \in B_1$ *and* $b \in \mathcal{U}^k$*.*

*Proof.* Note that $[j_0] = [j]$ in $KK(B_1, B_2)$. Let $\tau$ be the unique tracial state of $B_2$ then

$$\tau \circ j = \tau \circ j_0.$$

Since $K_1(B_1) = \{0\}$, $\tilde{\eta}_{j_0, j} = 0$. It follows theorem 9.1 of [49] (see also Theorem 1 of [51]) that (e 8.325) holds.

□

# 9 The general cases

In this section, we will show the main theorem of this paper. We will repeat the same argument deployed for the proof of 7.1 and use induction to present the proof.

**Definition 9.1.** Let $C$ be a unital separable amenable $C^*$-algebra in $\mathcal{N}$.

(1) We say $C$ has the property (P1) with integer $k \geq 1$, if the following statement holds:

Let $A \cong A \otimes \mathcal{U}$ be a unital simple AF-algebra with a unique tracial state $\tau$ and with $K_0(A) = \rho_A(K_0(A))$. There exists at least one unital monomorphism from $C \rtimes_\Lambda \mathbb{Z}^k$ into $A$.

Suppose that $\varphi_1, \varphi_2 : C \rtimes_\Lambda \mathbb{Z}^k \to A$ are two unital monomorphisms such that

$$\tau \circ \varphi_1 = \tau \circ \varphi_2 \text{ and } \tilde{\eta}(\varphi_1, \varphi_2) \;=\; 0. \tag{e 9.326}$$

Then, there exists a continuous path of unitaries $\{w_t : t \in [0, \infty)\} \subset U(B \otimes \mathcal{U}^k)$ such that

$$\lim_{t\to\infty} w_t^*(\psi_1(a) \otimes 1_{\mathcal{U}^k})w_t = \psi_2(a) \otimes 1_{\mathcal{U}^k} \text{ for all } a \in C \rtimes_\Lambda \mathbb{Z}^k, \tag{e 9.327}$$

where $A \subset B \cong B \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state, with $K_0(B) = \rho_B(K_0(B)$ which is also a subring of $\mathbb{R}$ and with $1_A = 1_B$, and where $\psi_i = j \circ \varphi_i^{(k)}$ (where $j : A \otimes \mathcal{U}^k \to B$ is a unital monomorphism ), $i = 1, 2.$ defined in 4.3) $i = 1, 2,$

(2) We say $C$ has the property (P2) with integer $k$, if the following statement holds:

Let $\Lambda : \mathbb{Z}^k \to Aut(C)$ be a homomorphism, let $A \cong A \otimes \mathcal{U}$ be a unital simple AF-algebra with a unique tracial state and with $K_0(A) = \rho_A(K_0(A))$ which is also a subring of $\mathbb{R}$. Suppose that there is at least one unital monomorphism from $C \rtimes_\Lambda \mathbb{Z}^k$ to $A$. Suppose that $j : C \rtimes_\Lambda \mathbb{Z}^k \to A$

is a unital monomorphism. Let $\epsilon > 0$, $\mathcal{F} \subset C \rtimes_\Lambda \mathbb{Z}^k$ be a finite subset and let $\mathcal{P} \subset K_1(C \rtimes_\Lambda \mathbb{Z}^k)$ be a finite subset. There is $\delta > 0$ and a finite subset $\{y_1, y_2, ..., y_m\} \subset K_1(C \rtimes_\Lambda \mathbb{Z}^k)$ satisfying the following:

If $\beta : K_1(C \rtimes_\Lambda \mathbb{Z}^k) \to K_0(A)$ is a homomorphism with

$$\rho_A((\beta(y_j))(\tau) < \delta, \;\; j = 1, 2, ..., m, \tag{e 9.328}$$

where $\tau$ is the unique tracial state of $A$, then there exists a unitary $u \in B$ such that

$$\|[\psi(a),\, u]\| < \epsilon \;\text{ for all }\; a \in \mathcal{F} \;\text{ and }\; \mathrm{bott}_1(\psi,\, u)|_{\mathcal{P}} = \beta'|_{\mathcal{P}} \tag{e 9.329}$$

where $\psi = j^{(k)}$ and $\beta' = (j_2)_{*0} \circ \beta$ ($j_2 : A$ is as in 4.3), where $A \subset B \cong B \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state, with $K_0(B) = \rho_B(K_0(B)$ which is also a subring of $\mathbb{R}$ and with $1_A = 1_B$ ($j_2 : A \to B$ is the unital embedding).

(3) We say that $C$ has the property (P3) with integer $k$, if the following statement holds:

Let $\Lambda : \mathbb{Z}^k \to Aut(C)$ be a homomorphism and let $A \cong A \otimes \mathcal{U}$ be a unital simple AF-algebra with a unique tracial state $\tau$ and $\rho_A(K_0(A)) = K_0(A)$. There exists at least one unital monomorphism from $C \rtimes_\Lambda \mathbb{Z}^k$ into $A$.

Let $h : (C \rtimes_\Lambda \mathbb{Z}^k) \otimes C(\mathbb{T}) \to A$ be a unital monomorphism. Let $\{\alpha_1, \alpha_2, ..., \alpha_k\}$ be the set of generators of $\Lambda(\mathbb{Z}^k)$ and defined $\Lambda_{k-1} : \mathbb{Z}^{k-1} \to Aut(A)$ by $\{\alpha_1, \alpha_2, ..., \alpha_{k-1}\}$. For any $\epsilon > 0$ and a finite subset $\mathcal{F} \subset C \rtimes_\Lambda \mathbb{Z}^k$, there exists $\delta > 0$, $\eta > 0$, a finite subset $\mathcal{G} \subset C \rtimes_\Lambda \mathbb{Z}^k$, a finite subset $\bar{\mathcal{G}} \subset C \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1} \otimes C(\mathbb{T})$ and a finite subset $\mathcal{P} \subset K_1(C \rtimes_\Lambda \mathbb{Z}^k)$ satisfying the following:

Suppose that there is a unitary $v \in U(A)$ and a contractive completely positive linear map $L : (C \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1}) \otimes C(\mathbb{T}) \to A$ such that

$$\|[h(a),\, v]\| < \delta \;\text{ for all }\; a \in \mathcal{G}, \mathrm{bott}_1(h, v)|_{\mathcal{P}} = 0 \;\text{ and } \tag{e 9.330}$$
$$\tau \circ h|_{(C \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1}) \otimes C(\mathbb{T})} \approx_\eta \tau \circ L \;\text{ on }\; \bar{\mathcal{G}}, \tag{e 9.331}$$

and

$$L \approx_\delta h \;\text{ on }\; \mathcal{G} \;\text{ and }\; L(1 \otimes z) \approx_\delta v \tag{e 9.332}$$

and where $z \in C(\mathbb{T})$ is the identity function on the unit circle. Then there exists a unitary $W \in B \otimes \mathcal{U}^k$ such that

$$\|[h^{(k)}(c),\, W]\| \;\; < \;\; \epsilon \;\text{ for all }\; c \in \mathcal{F} \;\text{ and } \tag{e 9.333}$$
$$W^*(v \otimes 1_{\mathcal{U}^k})W \;\; \approx_\epsilon \;\; h(1 \otimes z) \otimes 1_{\mathcal{U}^k}, \tag{e 9.334}$$

where $A \subset B \cong B \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state, with $K_0(B) = \rho_B(K_0(B)$ which is also a subring of $\mathbb{R}$ and with $1_A = 1_B$.

(4) We say that $C$ has the property (P4) with integer $k$, if the following holds:

Let $\Lambda : \mathbb{Z}^k \to Aut(C)$ be a homomorphism and let $A \cong A \otimes \mathcal{U}$ be a unital simple AF-algebra with unique tracial state $\tau$ and $\rho_A(K_0(A)) = K_0(A)$. There exists at least one unital monomorphism from $C \rtimes_\Lambda \mathbb{Z}^k$ into $A$. For any $\epsilon > 0$ and a finite subset $\mathcal{F} \subset C \rtimes_\Lambda \mathbb{Z}^k$, there exists $\delta > 0$, a finite subset $\mathcal{G} \subset C \rtimes_\alpha \mathbb{Z}$ and a finite subset $\mathcal{P} \subset \underline{K}(C \rtimes_\Lambda \mathbb{Z}^k)$ satisfying the following:

Suppose that $\varphi : C \rtimes_\Lambda \mathbb{Z}^k \to A$ is a unital monomorphism and suppose that there is a unitary $v \in U(A)$ such that

$$\|[\varphi(a),\, v]\| < \delta \;\text{ for all }\; a \in \mathcal{G} \;\text{ and }\; \mathrm{bott}_1(\varphi, v)|_{\mathcal{P}} = 0. \tag{e 9.335}$$

Then there exists a continuous path of unitaries $\{v_t : t \in [0,1]\} \subset U(B \otimes \mathcal{U}^k)$ such that

$$v_0 = v \otimes 1_{\mathcal{U}^k}, \;\; v_1 = 1_{A \otimes \mathcal{U}^k}, \;\; \|[\varphi^{(k)}(a), v_t]\| < \epsilon \tag{e 9.336}$$

for all $a \in \mathcal{F}$ and $t \in [0,1]$, and

$$\mathrm{Length}(\{v_t\}) \le 2\pi + \min\{\epsilon, 1\}., \tag{e 9.337}$$

where $A \subset B \cong B \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state, with $K_0(B) = \rho_B(K_0(B)$ which is also a subring of $\mathbb{R}$ and with $1_A = 1_B$.

**Lemma 9.2.** *Let $C_0$ be a unital separable amenable $C^*$-algebra in $\mathcal{N}$ which has the property* (P4) *with integer $k-1>0$ and let $\Lambda : Z^k \to Aut(C_0)$ be a homomorphism such that $\{\alpha_1, \alpha_2, ..., \alpha_k\}$ forms a set of generators of $\Lambda(\mathbb{Z}^k)$. Let $\Lambda_{k-1} : \mathbb{Z}^{k-1} \to Aut(C_0)$ be defined by $\{\alpha_1, \alpha_2, ..., \alpha_{k-1}\}$.*

*For any $\epsilon > 0$ and finite subset $\mathcal{F} \subset C_0 \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1}$, there is $\delta > 0$, a finite subset $\mathcal{G} \subset C_0 \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1}$ and a finite subset $\mathcal{P} \subset K_1(C_0 \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1})$ satisfying the following:*

*Suppose that $A \cong A \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state and with $K_0(A) = \rho_A(K_0(A))$, suppose that $\varphi : C_0 \rtimes_\Lambda \mathbb{Z}^k \to A$ is a unital monomorphism and suppose that there is a unitary $v \in U(A)$ such that*

$$\|[v, \varphi(a)]\| < \delta, \quad \textit{for all } a \in \mathcal{G} \textit{ and } \mathrm{bott}_1(\varphi, v)|_{\mathcal{P}} = 0 \tag{e 9.338}$$

*then there is a unitary $w \in B \otimes \mathcal{U}^k$ such that*

$$\begin{aligned} w^*(\varphi^{(k-1)}|_{C_0 \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1}}(a) \otimes 1_{\mathcal{U}^k})w \;\; &\approx_\epsilon \;\; \varphi^{(k-1)}|_{C_0 \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1}}(a) \otimes 1_{\mathcal{U}^k} && \text{(e 9.339)} \\ \textit{for all } a \in \mathcal{F} \textit{ and} & && \text{(e 9.340)} \\ w^* u (v \otimes 1_{\mathcal{U}^k}) w \;\; &\approx_\epsilon \;\; u, && \text{(e 9.341)} \end{aligned}$$

*where $u = \varphi^{(k)}(u_\alpha)$ and where $A \subset B \cong B \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state, with $K_0(B) = \rho_B(K_0(B)$ which is also a subring of $\mathbb{R}$ and with $1_A = 1_B$.*

*Proof.* Let $C = C_0 \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1}$. The proof is exactly the same as that of 3.2. However, instead of using $\varphi'$, we use $\varphi^{(k-1)}$ and instead of applying the Basic Homotopy Lemma for AH-algebras, we apply the property (P4) with integer $k-1$.

□

**Theorem 9.3.** *Let $C_0$ be a unital separable amenable $C^*$-algebra in $\mathcal{N}$ which has the property* (P1) *with integer $k-1$ and* (P4) *with integer $k-1>0$.*

*Let $\Lambda : \mathbb{Z}^k \to Aut(C)$ be a homomorphism and let $A \cong A \otimes \mathcal{U}$ be a unital simple AF-algebra with a unique tracial state $\tau$ and $K_0(A) = \rho_A(K_0(A))$. Suppose that $\{\alpha_1, \alpha_2, ..., \alpha_k\}$ forms a set of generators for $\Lambda(\mathbb{Z}^k)$. Denote $\Lambda_{k-1} : \mathbb{Z}^{k-1} \to Aut(C_0)$ be defined by $\{\alpha_1, \alpha_2, ..., \alpha_{k-1}\}$.*

*Suppose that $\varphi_1, \varphi_2 : C \rtimes_\Lambda \mathbb{Z}^k \to A$ are two unital monomorphisms such that*

$$\tau \circ \varphi_1 = \tau \circ \varphi_2. \tag{e 9.342}$$

*Suppose also that*

$$R_{\varphi_1 \circ j_0, \varphi_2 \circ j_0}(K_1(C_0 \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1})) \subset \rho_A(K_0(A)), \tag{e 9.343}$$

*where $j_0 : C_0 \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1} \to C_0 \rtimes_\Lambda \mathbb{Z}^k$ is the embedding.*

*Then there exists a sequence of unitaries $\{w_n\} \subset U(B \otimes \mathcal{U}^k)$ such that*

$$\lim_{n \to \infty} \mathrm{ad}\, w_n \circ \psi_1^{(k)}(a) = \psi_2^{(k)}(a) \;\; \textit{for all } a \in C \rtimes_\alpha \mathbb{Z}, \tag{e 9.344}$$

*where $A \subset B \cong B \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state, with $K_0(B) = \rho_B(K_0(B)$ which is also a subring of $\mathbb{R}$ and with $1_A = 1_B$.*

*Proof.* Let $C = C_0 \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1}$. The proof of this is exactly the same as that of 3.6. However, instead of applying 10.7 of [49], (where AH-algebra is used) we use the property (P1), and instead of applying 3.2, we apply 9.2.

$\square$

**Theorem 9.4.** *Let $C_{00}$ be a unital separable amenable $C^*$-algebra in $\mathcal{N}$ which has the property* (P1) *with integer $k-1$.*

*Then $C_{00}$ has the property* (P2) *with $k$.*

*Proof.* Let $\Lambda : \mathbb{Z}^k \to Aut(C_{00})$ be homomorphism, let $\{\alpha_1, \alpha_2, ..., \alpha_k\}$ forms a set of generators for $\Lambda(\mathbb{Z}^k)$ and let $\Lambda_{k-1} : \mathbb{Z}^{k-1} \to Aut(C_{00})$ defined by $\{\alpha_1, \alpha_2, ..., \alpha_{k-1}\}$. Define $C_0 = C_{00} \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1}$. The proof is very much the same of that 4.4 but we apply the properties (P1) instead of using the assumption that $C_0$ is an AH-algebra. We proceed the same proof until (e 4.137). Note that, since $C_{00}$ satisfies property (P1) with integer $k-1$, 9.3 holds. Next, instead of applying 3.6, we apply 9.3 to obtain a unital simple AF-algebra $B_1 \supset B$ ($B$ as in the proof of 4.4) with a unique tracial state, with $K_0(B_1) = \rho_B(K_0(B_1))$ which is also a subring of $\mathbb{R}$ and with $1_B = 1_{B_1}$ for which $B_1 \cong B_1 \otimes \mathcal{U}$, and a unitary $w \in B_1 \otimes \mathcal{U}^k$ such that (e 4.138)holds with $\psi_1$ replaced $(h \circ j_1)^{(k)}$ and $\psi$ replaced by $j^{(k)}$. Then (e 4.139) follows (with $\psi$ replaced by $j^{(k)}$).

$\square$

**Lemma 9.5.** *Let $C_{00}$ be a unital separable amenable $C^*$-algebra in $\mathcal{N}$ which has the property* (P2) *with $k-1$, let $\Lambda : \mathbb{Z}^k \to Aut(C_{00})$ be a homomorphism such that $\{\alpha_1, \alpha_2, ..., \alpha_k\}$ forms a set of generators of $\Lambda(\mathbb{Z}^k)$ and let $\Lambda_{k-1} : \mathbb{Z}^{k-1} \to Aut(C_{00})$ be defined $\{\alpha_1, \alpha_2, ..., \alpha_{k-1}\}$.*

*Let $C_0 = C_{00} \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1}$, let $A \cong A \otimes \mathcal{U}$ be a unital simple AF-algebra with a unique tracial state $\tau$ and with $K_0(A) = \rho_A(K_0(A))$ and let $j : C_{00} \rtimes_\Lambda \mathbb{Z}^k \to A$ be a unital monomorphism, let $\epsilon > 0$, $\mathcal{F} \subset C_0$ be a finite subset and $\mathcal{P} \subset \underline{K}(C_0)$ be a finite subset. There is $\delta > 0$, a finite subset $\mathcal{G} \subset C_0$, and integer $K$ and a finite subset $\mathcal{Q} \subset \underline{K}(C_{00} \rtimes_\Lambda \mathbb{Z}^k)$ satisfying the following:*

*Suppose that $v \in U(A)$ is a unitary such that*

$$\|[j(a),\, v]\| < \delta \text{ for all } a \in \mathcal{G} \text{ and} \tag{e 9.345}$$

$$[L]|_{\mathcal{Q}} = [j]|_{\mathcal{Q}}, \tag{e 9.346}$$

*where $L : C_{00} \rtimes_\Lambda \mathbb{Z}^k \to A$ is a contractive completely positive linear map such that*

$$\|L(\sum_{k=-K}^{K} f_k u_\alpha^k) - \sum_{k=-K}^{K} j(f_k)(j(u_\alpha)v)^k\| < \delta \tag{e 9.347}$$

*for all $f_k \in \mathcal{G}$. Then there exists a unitary $w \in U(B \otimes \mathcal{U}^k)$ such that*

$$\|[j^{(k)}(a),\, w]\| < \epsilon \text{ for all } a \in \mathcal{F} \text{ and} \tag{e 9.348}$$

$$\text{bott}_1(j^{(k)}|_{C_0},\, j^{(k)}(u_\alpha)^* w^* j^{(k)}(u_\alpha)(v \otimes 1_{\mathcal{U}^k})w)|_{\mathcal{P}} = 0, \tag{e 9.349}$$

*where $A \subset B \cong B \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state, with $K_0(B) = \rho_B(K_0(B))$ which is also a subring of $\mathbb{R}$ and with $1_A = 1_B$.*

*Proof.* The proof of this lemma is almost the same as that of 5.2 but we apply the property (P2) (with $k-1$) instead of Lemma 7.3 of [49]. We proceed as follows.

We will keep the first paragraph of that proof and keep the line which defines $\mathcal{P}$. Let $\eta > 0$ (in place of $\delta$), $\{y_1, y_2, ..., y_m\} \subset K_1(C_0))$ be required by (P2) for $\epsilon_0/2 > 0$, $\{x_1, x_2, ..., x_k\}$ and

$\mathcal{F}_1$. We then remove the paragraph related to Lemma 7.3 of [49] right after the line which defines $\mathcal{P}$.

We then continue the same proof until the line right after (e 5.162). We replace the next line by the following:

By (e 5.149) and by property (P2) with $k-1$ for $C_{00}$, there is a unitary $w \in B \otimes \mathcal{U}^k$ such that (e 5.163) holds with $j$ replaced by $j^{(k)}$, where $A \subset B \cong B \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state, with $K_0(B) = \rho_B(K_0(B)$ which is also a subring of $\mathbb{R}$ and with $1_A = 1_B$. The rest of the proof remains the same by replacing $j$ by $j^{(k)}$ and $v$ by $v \otimes 1_{\mathcal{U}^k}$.

□

**Theorem 9.6.** *Let $C_0$ be a unital separable amenable $C^*$-algebra in $\mathcal{N}$ which has property* (P1)*,* (P2)*,* (P3) *and* (P4) *with $k-1$.*

*Then $C_0$ has the property* (P3) *with integer $k$ .*

*Proof.* The proof is the same as that of 5.3. The main difference is that we will use property (P3) instead of 2.13.

We proceed as follows:

Let $\Lambda : \mathbb{Z}^k \to Aut(C_0)$ be a monomorphism such that $\{\alpha_1, \alpha_2, ..., \alpha_k\}$ forms a set of generators for $\Lambda(\mathbb{Z}^k)$ and let $\Lambda_j : \mathbb{Z}^j \to Aut(C_0)$ defined by $\{\alpha_1, \alpha_2, ..., \alpha_j\}$ $(1 \le j \le k-1)$.

Define $C = C_0 \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1}$.

Since $C_0$ has property (P1), (P2), (P3) and (P4) (with $k-1$), 9.2 and 9.5 hold.

We will not make changes from the beginning of the proof to (e 5.177) and keep the definition of $\tau_0$ with only a couple of exceptions: We will not apply 3.2, but apply 9.2, not 5.2 but 9.5. So we replace 3.2 by 9.2 and replace 5.2 by 9.5 in these lines. We also need, however, replace $\alpha$ by $\alpha_k$.

We may also assume that $\mathcal{G}_2 = \mathcal{G}_0 \cup \{1 \otimes z\}$ for some finite subset $\mathcal{G}_0 \subset C$ and $\mathcal{G}_2' = \mathcal{G}_0' \cup \{1 \otimes z\}$ for some finite subset $\mathcal{G}_0'$.

After we define $\tau_0$ (right after (e 5.177)), we will apply property (P3) with integer $k-1$.

Let $0 < \delta_3 < \delta_2$ (in place of $\delta$), $\eta > 0$, $\tilde{\mathcal{G}} \subset C_0 \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1}$ (in place of $\mathcal{G}$) be a finite subset, $\bar{\mathcal{G}}_1 \subset (C_0 \rtimes_{\Lambda_{k-2}} \mathbb{Z}^{k-2}) \otimes C(\mathbb{T})$ (in place of $\bar{\mathcal{G}}$) be a finite subset required and $\mathcal{P}_4 \subset K_1(C)$ be a finite subset required by (P3) (with $k-1$) for $\min\{\delta_2/4, \epsilon/4\}$ and $\mathcal{G}_0'$.

As in the proof of 5.3, without loss of generality, we may assume that $\bar{\mathcal{G}}_1 = \mathcal{G}_3' \otimes \mathcal{G}_3''$, where $\{1_C\} \subset \mathcal{G}_3' \subset C$ and $\{1_{C(\mathbb{T})}, z\} \subset \mathcal{G}_3'' \subset C(\mathbb{T})$ are finite subsets.

We delete the lines after the definition of $\tau_0$ until the definition of $j_0$ of the proof of 5.3. Keep the definition of $j_0$ and put $\mathcal{P} = (j_0)_{*1}(\mathcal{P}_4)$ and keep the same $\eta_1$. We keep the next few lines but stop right after (e 5.180) and replace (e 5.172) by (e 9.330), (e 5.173) by (e 9.331) and (e 5.174) by (e 9.332). Since we assume that (e 9.332) holds, we have

$$\text{bott}_1(h|_C,\, v)|_{\mathcal{P}_4} = 0. \tag{e 9.350}$$

By applying property (P3) (with $k-1$), we obtain a unitary $w_0' \in B_1 \otimes \mathcal{U}^{k-1}$ such that

$$\begin{aligned} \|[(h|_C)^{(k-1)}(c),\, w_0']\| &< \min\{\delta_2/4, \epsilon/2\} \text{ for all } c \in \mathcal{G}_0' \text{ and} & \text{(e 9.351)}\\ (w_0')^*(v \otimes 1_{\mathcal{U}^{k-1}})w_0' &\approx_{\min\{\delta_2/4,\epsilon/2\}} h(1_{C_0} \otimes z) \otimes 1_{\mathcal{U}^{k-1}}, & \text{(e 9.352)} \end{aligned}$$

where $A \subset B_1 \cong B_1 \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state, with $K_0(B_1) = \rho_{B_1}(K_0(B_1))$ which is also a subring of $\mathbb{R}$ and with $1_A = 1_{B_1}$.

Define $w_0 = w_0' \otimes 1_{\mathcal{U}}$. Then

$$\begin{aligned} \|[h^{(k)}(c),\, w_0]\| &< \min\{\delta_2/4, \epsilon/2\} \text{ for all } c \in \mathcal{G}_0' \text{ and} & \text{(e 9.353)}\\ (w_0)^*(v \otimes 1_{\mathcal{U}^k})w_0 &\approx_{\min\{\delta_2/4,\epsilon/2\}} h(1_{C_0} \otimes z) \otimes 1_{\mathcal{U}^k} & \text{(e 9.354)} \end{aligned}$$

Define $V = (h^{(k)}(u_{\alpha_k}))^* w_0^* h^{(k)}(u_{\alpha_k}) w_0$. We then continue the same proof as in 5.3 as follows. We have

$$\|[h^{(k)}(a),\, V]\| < \min\{\epsilon/2, \delta_2/4\} \ \text{ for all } \ a \in \mathcal{G}_2'. \tag{e 9.355}$$

Thus, there exists a unital contractive completely positive linear map $L_1 : C \rtimes_\Lambda \mathbb{Z}^k \otimes C(\mathbb{T}) \to A \otimes \mathcal{U}^k$ which satisfies (e 5.176) (replacing $\alpha$ by $\alpha_k$ and $h$ by $h^{(k)}$) with $v' = V$. Therefore

$$L_1 \approx_{\delta_2} h^{(k)} \ \text{ on } \ \mathcal{G}_2'' \ \text{ and } \ L_1(u_{\alpha_k}) \approx_{\delta_2} h^{(k)}(u_{\alpha_k})V. \tag{e 9.356}$$

Note that

$$[L_1]|_{\mathcal{P}_2} = [h^{(k)}]|_{\mathcal{P}_2} \tag{e 9.357}$$

By the assumption (e 9.330) and by (e 9.354),

$$\begin{aligned} \mathrm{bott}_1(\mathrm{ad}\, w_0 \circ h^{(k)},\, v \otimes 1_{\mathcal{U}^k}]]\|\|_{\mathcal{P}_3} &= \mathrm{bott}_1(\mathrm{ad}\, w_0 \circ h^{(k)},\, w_0^*(v \otimes 1_{\mathcal{U}^k})w_0)|_{\mathcal{P}_3} && \text{(e 9.358)} \\ &= \mathrm{bott}_1(h^{(k)},\, v)|_{\mathcal{P}_3} = 0. && \text{(e 9.359)} \end{aligned}$$

It follows that (since $\ker\rho_{B_1} = \{0\}$)

$$[L_1]|_{\boldsymbol{\beta}(\mathcal{P}_3)} = 0 = [h^{(k)}]|_{\boldsymbol{\beta}(\mathcal{P}_3)}. \tag{e 9.360}$$

Therefore, by applying 9.5, there is a unitary $w_1 \in U((B_2 \otimes \mathcal{U}^k)$ such that

$$\begin{aligned} \|([h^{(k)})^{(k)}(a),\, w_1]\| &< \delta_1/2 \ \text{ for all } \ a \in \mathcal{G}_1 \ \text{ and} && \text{(e 9.361)} \\ \mathrm{bott}_1((h^{(k)})^{(k)},\, V_1)|_{\mathcal{P}_1} = 0, & && \text{(e 9.362)} \end{aligned}$$

where $V_1 = (h^{(k)})^{(k)}(u_{\alpha_k})^* w_1^* (h^{(k)})^{(k)}(u_{\alpha_k})(V \otimes 1_{\mathcal{U}^k})w_1$ and where $B_1 \otimes \mathcal{U}^k \subset B_2 \cong B_2 \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state, with $K_0(B_2) = \rho_{B_2}(K_0(B_2))$ which is also a subring of $\mathbb{R}$ and with $1_{B_1 \otimes \mathcal{U}^k} = 1_{B_2}$. Note that

$$(h^{(k)})^{(k)}(u_{\alpha_k})(V \otimes 1_{\mathcal{U}^k}) = w_0^* h^{(k)}(u_{\alpha_k}) w_0 \otimes 1_{\mathcal{U}^{k-1}} \otimes \imath(u_\sigma).$$

It follows from 9.2 that there exists a unitary $w_2 \in U(B_2 \otimes \mathcal{U}^{2k})$ such that

$$\begin{aligned} w_2^*(((h^{(k)})^{(k)})^{(k)}(c))w_2 &\approx_{\epsilon/2} ((h^{(k)})^{(k)})^{(k)}(a) \ \text{ for all } \ a \in \mathcal{F} && \text{(e 9.363)} \\ w_2^*(w_1^*(w_0^* h^{(k)}(u_{\alpha_k}) w_0 \otimes 1_{\mathcal{U}^k})w_1) \otimes 1_{\mathcal{U}^k} \otimes \imath(u_{\alpha_k}) &\approx_{\epsilon/2} (h^{(k)})^{(k)}(u_{\alpha_k}) \otimes 1_{\mathcal{U}^{k-1}} \otimes \imath(u_{\alpha_k}). && \text{(e 9.364)} \end{aligned}$$

Define $W_1 = ((w_0 \otimes 1_{\mathcal{U}})w_1) \otimes 1_{\mathcal{U}})w_2$. Then, by (e 9.353) and (e 9.354),

$$\begin{aligned} \|[((h^{(k)})^{(k)})^{(k)}(c),\, W_1]\| &< \epsilon \ \text{ for all } \ c \in \mathcal{F} \ \text{ and} && \text{(e 9.365)} \\ W_1^*(v \otimes 1_{\mathcal{U}^{3k}})W &\approx_\epsilon h(1 \otimes z) \otimes 1_{\mathcal{U}^{3k}}. && \text{(e 9.366)} \end{aligned}$$

Finally, we apply the absorption lemma 8.5 twice and 8.7 once.

□

**Lemma 9.7.** *Let $C$ be a unital separable amenable $C^*$-algebra in $\mathcal{N}$ and let $\Lambda : \mathbb{Z}^k \to Aut(C)$ be a homomorphism. Suppose that $C$ has property (P3) for $k$. Then $C$ has the property* (P4) *with $k$.*

*Proof.* The proof follows from that of 5.6 almost verbatim. We replace the application of 5.3 by the application of property (P3). So in (e 5.227), $W$ should be in $U(B_1 \otimes \mathcal{U}^k)$, $\bar{v}$ should be replaced by $\bar{v} \otimes 1_{\mathcal{U}^k}$, $h(1 \otimes z) \otimes 1_{\mathcal{U}}$ replaced by $h(1 \otimes z) \otimes 1_{\mathcal{U}^k}$ and $\psi$ should be replaced by $\varphi^{(k)}$, where $A \otimes \mathcal{U} \subset B \cong B \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state, with $K_0(B) = \rho_B(K_0(B))$ which is also a subring of $\mathbb{R}$ and with $1_{A\otimes\mathcal{U}} = 1_B$. For the rest of the proof, we will continue use this $W$ and replace $1_{\mathcal{U}}$ by $1_{\mathcal{U}^k}$, $\psi$ by $\varphi^{(k)}$ and $V$ by $V \otimes 1_{\mathcal{U}^k}$ (and $\lambda(t)$ and $v_t$ are in $B \otimes \mathcal{U}^k$).

□

**Theorem 9.8.** *Let $C$ be a unital separable amenable $C^*$-algebra in $\mathcal{N}$ and let $\Lambda : \mathbb{Z}^k \to Aut(C)$ be a monomorphism.*

*Suppose that $C$ satisfies property* (P1)*,* (P2)*,* (P3) *and* (P4) *for $k-1$. Then $C$ has property* (P1)*,* (P2)*,* (P3) *and* (P4) *for $k$.*

*Proof.* Since $C$ has property (P1) (with integer $k-1$), 9.3 holds. It follows from 9.4 and 9.6 that $C$ has the property (P2) and (P3) for $k$. By 9.7, $C$ also has the property (P4) for $k$. It remains to show that $C$ has property (P1) for $k$.

The proof is exactly the same as that of 6.1.

Suppose that $\{\alpha_1, \alpha_2, ..., \alpha_k\}$ forms a set of generators for $\Lambda(\mathbb{Z}^k)$. Let $\Lambda_{k-1} : \mathbb{Z}^{k-1} \to Aut(C)$ be defined by $\{\alpha_1, \alpha_2, ..., \alpha_{k-1}$.

In the proof of 6.1, we need to replace $C$ by $C \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1}$, $\alpha$ by $\alpha_k$, $\psi_1$ and $\psi_2$ by $\varphi_1^{(k)}$ and $\varphi_2^{(k)}$, respectively. In particular, (e 6.248) holds for $\varphi_1^{(k)}$ and $\varphi_2^{(k)}$ (instead of $\psi_1$ and $\psi_2$). We also replace $A \otimes \mathcal{U}$ by $B_1 \otimes \mathcal{U}^k$ for some unital simple AF-algebra $B_1$ which has the properties: (1) $A \subset B_1 \cong B_1 \otimes \mathcal{U}$, (2) $K_0(B_1) = \rho_{B_1}(K_0(B_1))$ is a subring of $\mathbb{R}$, (3) $1_A = 1_{B_1}$.

We will prepare to apply (P4) (with $k$) instead of 5.6. We will apply property (P2) (with $k$). Hence (e 6.258) holds by replacing $\psi_2'$ by $(\varphi_2^{(k)})^{(k)}$. We proceed the same calculation and apply property (P4) (with $k$) to obtain (e 6.281) with $\psi_2''$ being replaced by $((\varphi_2^{(k)})^{(k)})^{(k)}$ and with $v_n(t) \in B_2 \otimes \mathcal{U}^k$, for some unital simple AF-algebra $B_2 \cong B_2 \otimes \mathcal{U}$ which satisfies: (1) $B_1 \otimes \mathcal{U}^k \subset B_2$, (2) $K_0(B_2) = \rho_{B_2}(K_0(B_2))$ which is a subring of $\mathbb{R}$, (3) $1_{B_1\otimes\mathcal{U}^k} = 1_{B_2}$.

We then obtain (e 6.282) with $\psi_1''$ and $\psi_2''$ being replaced by $(\varphi_1^{(k)})^{(k)})^{(k)}$ and $(\varphi_2^{(k)})^{(k)})^{(k)}$, and with $u(t) \in B_2 \otimes \mathcal{U}^{2k}$.

Finally, to obtain (e 6.241), we apply the absorption lemma 8.5 twice.

□

**Theorem 9.9.** *Let $C$ be a unital separable amenable $C^*$-algebra in $\mathcal{N}$. Suppose that $\Lambda : \mathbb{Z}^k \to Aut(C)$ is a homomorphism so that $\{\alpha_1, \alpha_2, ..., \alpha_k\}$ forms a set of generators for $\Lambda(\mathbb{Z}^k)$ and suppose that $\Lambda_{k-1} : \mathbb{Z}^{k-1} \to Aut(C)$ defined by $\{\alpha_1, \alpha_2, ..., \alpha_{k-1}\}$.*

*Suppose that there is a unital monomorphism $\varphi : C \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1} \to A$, where $A \cong A \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state $\tau$ and with $K_0(A) = \rho_A(K_0(A))$ such that $\tau \circ \varphi$ is $\alpha_k$-invariant.*

*Suppose $C$ has property* (P1) *with $k-1$. Then there is a unital monomorphism $\psi : C \rtimes_\Lambda \mathbb{Z}^k \to B$ for some unital simple AF-algebra with a unique tracial state $\tau_1$ and with $K_0(B) = \rho_B(K_0(B))$ such that*

$$\tau_1 \circ \psi \circ j_0 = \tau \circ \varphi, \tag{e 9.367}$$

*where $j_0 : C \rtimes_{\Lambda_{k-1}} \mathbb{Z}^{k-1} \to C \rtimes_\Lambda \mathbb{Z}^k$ is the embedding.*

*Proof.* The proof is an application of Theorem 5.5 of [49] exactly as in the proof of 7.1 (see also 7.2).

□

**Theorem 9.10.** *Let $C$ be a unital separable $AH$-algebra and let $\Lambda : \mathbb{Z}^k \to Aut(C)$ be a monomorphism. Then $C$ can be embedded into a unital simple $AF$-algebra if and only if $C$ admits a faithful $\Lambda$-invariant tracial state.*

*Proof.* It suffices to show the "if part" of the theorem.

Suppose that $\{\alpha_1, \alpha_2, ..., \alpha_k\}$ forms a set of generators for $\Lambda(\mathbb{Z}^k)$. Denote by $\Lambda_j : Z^j \to Aut(C)$ defined by $\{\alpha_1, \alpha_2, ..., \alpha_j\}$, $j = 1, 2, ..., k$.

By 4.4, 5.3, 5.6 and 6.1, $C$ has property (P1) (P2), (P3), (P4) for $k = 1$. Theorem 9.9 (or 8.6) implies that there is a unital monomorphism $\varphi_2 : C \rtimes_{\Lambda_2} \mathbb{Z}^2 \to A_2$, where $A_2 \cong A_2 \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state $\tau_2$ and with $K_0(A_2) = \rho_{A_2}(K_0(A_2))$ such that

$$\tau_2 \circ \varphi_2 \circ j_1 = t, \qquad (e\,9.368)$$

where $t$ is the chosen faithful $\Lambda$-invariant tracial state on $C$. In particular,

$$\tau_2 \circ \varphi_2 = \tau_2 \circ \varphi_2 \circ \alpha_i, \;\; i = 1, 2, ..., k. \qquad (e\,9.369)$$

We use the induction on $k$ to prove the theorem.

Suppose that we have shown the following:

(1) $C$ has the property (P1), (P2), (P3) and (P4) for $i - 1$.

(2) There is a monomorphism $\varphi_{i-1} : C \rtimes_{\Lambda_{i-1}} Z^{i-1} \to A_{i-1}$, where $A_{i-1} \cong A_{i-1} \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state $\tau_{i-1}$ and with $K_0(A_{i-1}) = \rho_{A_{i-1}}(K_0(A_{i-1}))$ such that

$$\tau_{i-1} \circ \varphi_{i-1} = \tau_{i-1} \circ \varphi \circ \alpha_l, \;\; l = 1, 2, ..., k. \qquad (e\,9.370)$$

It follows from 9.8 that $C$ has property (P1), (P2),(P3) and (P4) for $i$. It follows from 9.9 that there is a unital monomorphism $\varphi_i : C \rtimes_{\Lambda_i} \mathbb{Z}^i \to A_i \cong A_i \otimes \mathcal{U}$ for some unital simple AF-algebra $A_i$ with a unique tracial state $\tau_i$ and with $K_0(A_i) = \rho_{A_i}(K_0(A_i))$ such that

$$\tau_i \circ \varphi_i \circ j_i = \tau_{i-1} \circ \varphi_{i-1}, \qquad (e\,9.371)$$

where $j_i : C \rtimes_{\Lambda_{i-1}} \mathbb{Z}^{i-1} \to C \rtimes_{\Lambda_i} \mathbb{Z}^i$ is the embedding. By (e 9.370), we have

$$\tau_i \circ \varphi_i = \tau_i \circ \varphi_i \circ \alpha_l, \;\; l = 1, 2, ...., k. \qquad (e\,9.372)$$

Thus, there is a unital monomorphism $\varphi_k : C \rtimes_\Lambda \mathbb{Z}^k \to A_k$, where $A_k \cong A_k \otimes \mathcal{U}$ is a unital simple AF-algebra with a unique tracial state $\tau$ and with $K_0(A_k) = \rho_{A_k}(K_0(A_k))$.

□

**Theorem 9.11.** *Let $C$ be a unital AH-algebra and let $G$ be a finitely generated (discrete)abelian group. Suppose that $\Lambda : G \to Aut(C)$ is a homomorphism. Then $C \rtimes_\Lambda G$ can be embedded into a unital simple AF-algebra if and only if $C$ admits a faithful $\Lambda$-invariant tracial state.*

*Proof.* We use an argument of Nate Brown (see Remark 11.10 of [10]). Write $G = \mathbb{Z}^k \oplus G_0$, where $G_0$ is a finite group. By the Green's theorem (Cor. 2.8 of [25]), there is a monomorphism $h : C \rtimes_\Lambda G \to (C \rtimes_{\Lambda|_{\mathbb{Z}^k}} \mathbb{Z}^k) \otimes \mathcal{K}$. Since $C \rtimes_{\Lambda|_{\mathbb{Z}^k}} \mathbb{Z}^k$ has a faithful $\Lambda|_{\mathbb{Z}^k}$ -invariant tracial state, there is a unital monomorphism $\varphi : C \rtimes_{\Lambda|_{\mathbb{Z}^k}} \mathbb{Z}^k \to A$ for some unital simple AF-algebra. Then $\varphi$ gives a monomorphism $\varphi_1 : (C \rtimes_{\Lambda|_{\mathbb{Z}^k}} \mathbb{Z}^k) \otimes \mathcal{K} \to A \otimes \mathcal{K}$. Thus $\varphi_1 \circ h$ gives a monomorphism from $C \rtimes_\Lambda G$ into $A \otimes \mathcal{K}$. Let $e = \varphi_1 \circ h(1)$. Then $e$ is a projection. Note that $e(A \otimes \mathcal{K})e$ is a unital simple AF-algebra.

□

**Corollary 9.12.** *Let $X$ be a compact metric space and $\Lambda : \mathbb{Z}^k \to Homeo(X)$ be a homomorphism, where $Homeo(X)$ is the group of all homeomorphisms on $X$. Then the crossed product $C(X) \rtimes_\Lambda \mathbb{Z}^k$ can be embedded into a unital simple AF-algebra if and only if there is a strictly positive $\Lambda$-invariant probability Borel measure.*

# References

[1] B. Blackadar and D. Handelman, *Dimension functions and traces on $C^*$-algebras* J. Funct. Anal. **45** (1982), 297–340.

[2] O. Bratteli, G. A. Elliott, D. Evans and A. Kishimoto, *Homotopy of a pair of approximately commuting unitaries in a simple $C^*$-algebra*, J. Funct. Anal. **160** (1998), 466–523.

[3] L. G. Brown, R. G. Douglas and P.A. Fillmore, *Unitary equivalence modulo the compact operators and extensions of $C^*$-algebras* Proceedings of a Conference on Operator Theory (Dalhousie Univ., Halifax, N.S., 1973), pp. 58–128. Lecture Notes in Math., Vol. 345, Springer, Berlin, 1973.

[4] L. G. Brown, R. G. Douglas and P. A. Fillmore *Extensions of $C^*$-algebras and $K$-homology.* Ann. of Math. **105** (1977), 265–324.

[5] B. Blackadar and E. Kirchberg *Generalized inductive limits of finite-dimensional $C^*$-algebras*, Math. Ann. **307** (1997), 343–380.

[6] B. Blackadar and E. Kirchberg *Inner quasidiagonality and strong NF algebras*, Pacific J. Math. **198** (2001), 307–329.

[7] L. G. Brown and M. Dădărlat, *Extensions of $C^*$-algebras and quasidiagonality*, J. London Math. Soc. **53** (1996), 582–600.

[8] N. Brown, *AF embeddability of crossed products of AF algebras by the integers*, J. Funct. Anal. **160** (1998), no. 1, 150–175.

[9] N. Brown, *Crossed products of UHF algebras by some amenable groups*, Hokkaido Math. J. **29** (2000), 201–211

[10] N. Brown, *On quasidiagonal $C^*$-algebras. Operator algebras and applications* 19–64, Adv. Stud. Pure Math., 38, Math. Soc. Japan, Tokyo, 2004.

[11] N. Brown and M. Dadarlat, *Extensions of quasidiagonal $C^*$-algebras and $K$-theory*, Operator algebras and applications, 65–84, Adv. Stud. Pure Math., 38, Math. Soc. Japan, Tokyo, 2004.

[12] M.D. Choi and E. Effros, *The completely positive lifting problem for $C^*$-algebras*, Ann. of Math. **104** (1976), 309-322.

[13] M. Dădărlat, *Quasidiagonal morphisms and homotopy*, J. Funct. Anal. 151 (1997), 213-

[14] M. Dădărlat, *On the approximation of quasidiagonal $C^*$-algebras*, J. Funct. Anal. **167** (1999), 69–78.

[15] M. Dădărlat, *Residually finite dimensional $C^*$-algebras and subquotients of the CAR algebra*, Math. Res. Lett. **8** (2001), 545–555.

[16] M. Dadarlat and T. Loring, *A universal multicoefficient theorem for the Kasparov groups*, Duke Math. J. **84** (1996), 355–377.

[17] K. Davidson, D. Herrero and N. Salinas, *Quasidiagonal operators, approximation, and $C^*$-algebras*, Indiana Univ. Math. J. **38** (1989), 973–998.

[18] S. Eilers, T. Loring and G. K. Pedersen, *Quasidiagonal extensions and AF algebras*, Math. Ann. **311** (1998), 233–249.

[19] G. A. Elliott, *On the classification of $C^*$-algebras of real rank zero*, J. Reine Angew. Math. **443** (1993), 179–219.

[20] G. A. Elliott and G. Gong, *On the classification of $C^*$-algebras of real rank zero. II*, Ann. of Math. **144** (1996), 497–610.

[21] G. A. Elliott, G. Gong and L. Li, *Injectivity of the connecting maps in AH inductive limit systems*, Canad. Math. Bull. **48** (2005), 50–68.

[22] G. A. Elliott and M. Rørdam, *Classification of certain infinite simple $C^*$-algebras, II*, Comment. Math. Hel **70** (1995), 615-638.

[23] R. Exel, *The soft torus and applications to almots commuting matrics*, Pacific J. Math., **160** (1993), 207-217.

[24] G. Gong and H. Lin, *Almost multiplicative morphisms and $K$-theory.* Internat. J. Math. **11** (2000), 983–1000.

[25] P. Green, *The structure of imprimitivity algebras*, J. Funct. Anay. **36** (1980), 88-104.

[26] D. Hadwin, *Strongly quasidiagonal $C^*$-algebras*, J. Operator Theory **18** (1987), 3–18.

[27] P. R. Halmos, *Quasitriangular operators*, Acta, Sci. Math. (Szeged) **29** (1968), 283-293.

[28] S. Hu, H. Lin and Y. Xue, *Limits of homomorphisms with finite-dimensional range*, Internat. J. Math. **16** (2005), 807–821.

[29] A. Kishimoto and A. Kumjian, *The* Ext *class of an approximately inner automorphism*, Trans. Amer. Math. Soc. **350** (1998), 4127–4148.

[30] A. Kishimoto and A. Kumjian, *The Ext class of an approximately inner automorphism. II*, J. Operator Theory **46** (2001), 99–122.

[31] H. Lin, *Exponential rank of $C^*$-algebras with real rank zero and the Brown-Pedersen conjectures*, J. Funct. Anal. **114** (1993), 1–11.

[32] L. Li, *Classification of simple $C^*$-algebras: inductive limits of matrix algebras over one-dimensional spaces*, J. Funct. Anal. **192** (2002), 1–51.

[33] H. Lin, *Tracially AF $C^*$-algebras*, Trans. Amer. Math. Soc. **353** (2001), 693–722.

[34] H. Lin, *Tracial topological ranks of $C^*$-algebras*, Proc. London Math. Soc., **83** (2001), 199-234.

[35] H. Lin, *An introduction to the classification of amenable $C^*$-algebras*, World Scientific Publishing Co., Inc., River Edge, NJ, 2001. xii+320 pp. ISBN: 981-02-4680-3.

[36] H. Lin, *Embedding an AH-algebra into a simple $C^*$-algebra with prescribed $KK$-data.* $K$-Theory **24** (2001), 135–156.

[37] H. Lin, *Residually finite dimensional and AF-embeddable $C^*$-algebras*, Proc. Amer. Math. Soc. **129** (2001), 1689–1696

[38] H. Lin, *Classification of simple $C^*$-algebras and higher dimensional noncommutative tori*, Ann. of Math. (2) 157 (2003), no. 2, 521–544.

[39] H. Lin, *Simple AH-algebras of real rank zero*, Proc. Amer. Math. Soc. **131** (2003), 3813–3819.

[40] H. Lin, *Classification of simple $C^*$-algebras with tracial topological rank zero*, Duke Math. J.,**125** (2004), 91-119.

[41] H. Lin, *Traces and simple $C^*$-algebras with tracial topological rank zero*, J. Reine Angew. Math. 568 (2004), 99–137.

[42] H. Lin, *Minimal Homeomorphisms and Approximate Conjugacy in Measure*, preprint (arXiv/math.OA/0501262).

[43] H. Lin, *An approximate universal coefficient theorem*, Trans. Amer. Math. Soc. **357** (2005), 3375–3405.

[44] H. Lin *Classification of homomorphisms and dynamical systems*, Trans. Amer. Math. Soc. **359** (2007), 859-895.

[45] H. Lin, *Unitary equivalences for essential extensions of $C^*$-algebras*, preprint arxiv.org/math.OA/0403236.

[46] H. Lin, *Embedding crossed products into a unital simple AF-algebra*, preprint, arxiv,org/ OA/0604047

[47] H. Lin, *Approximate homotopy of homomorphisms from $C(X)$ into a simple $C^*$-algebra*, preprint, arxiv.org/ OA/0612125.

[48] H. Lin, *AF-embedding of crossed products of AH-algebras by $\mathbb{Z}$ and asymptotic AF-embedding*, Indiana J. Math., accepted, arxiv.org/ OA/0612529.

[49] H. Lin, *Asymptotic unitary equivalence and asymptotic inner automorphism*, preprint arxiv.org/OA/0703610.

[50] T. Loring, *$K$-theory and asymptotically commuting matrices*, Canad. J. Math. **40** (1988), 197–216.

[51] H. Matui, *AF embeddability of crossed products of AT algebras by the integers and its application*, J. Funct. Anal. **192** (2002), 562–580.

[52] C. Pasnicu and N. C. Phillips *Crossed products by Z with Rokhlin property* , in preparation.

[53] M. Pimsner, *Embedding some transformation group $C^*$-algebras into AF-algebras*, Ergodic Theory Dynam. Systems **3** (1983), 613–626.

[54] S. Popa, *On local finite-dimensional approximation of $C^*$-algebras.* Pacific J. Math. **181** (1997), 141–158.

[55] M. Rørdam, *On the structure of simple $C^*$-algebras tensored with a UHF-algebra. II* J. Funct. Anal. **107** (1992), 255–269.

[56] J. Rosenberg, *Appendix to Strongly quasidiagonal $C^*$-algebras*, J. Operator Theory **18** (1987), 3-18.

[57] N. Salinas, *Relative quasidiagonality and $KK$-theory*, Houston J. Math., **18** (1992), 97-116.

[58] F. J. Thayer, *Quasidiagonal $C^*$-algebras*, J. Funct. Anal. **25** (1977), 50-57.

[59] J. Rosenberg and C. Schochet, *The Knneth theorem and the universal coefficient theorem for Kasparov's generalized $K$-functor*, Duke Math. J. **55** (1987), 431–474.

[60] A. Toms *On the classification problem for nuclear C*-algebras*, preprint, math.OA/0509103.

[61] J. Villadsen, *On the stable rank of simple $C^*$-algebras*, J. Amer. Math. Soc. **12** (1999) 1091–1102.

[62] D. Voiculescu, *A non-commutative Wely-con Nuemann theorem*, Rev. Roumanine Math., Pures Apply. **21** (1976), 97-113.

[63] D. Voiculescu, *Almost inductive limit automorphisms and embeddings into AF-algebras*, Ergodic Theory Dynam. Systems, **6** (1986), 475-484.

[64] D. Voiculescu, *A note on quasi-diagonal $C^*$-algebras and homotopy*, Duke Math. J. **62** (1991), 267–271.

[65] D. Voiculescu, *Around quasidiagonal operators* Integral Equations Operator Theory **17** (1993), 137–149.

[66] S. Zhang, *$K_1$-groups, quasidiagonality, and interpolation by multiplier projections*, Trans. Amer. Math. Soc. **325** (1991), 793–818.